

## CATALAN DETERMINANTS — A COMBINATORIAL APPROACH

ARTHUR T. BENJAMIN, NAIOMI T. CAMERON, JENNIFER J. QUINN, AND CARL R.  
YERGER

ABSTRACT. Determinants of matrices involving the Catalan sequence have appeared throughout the literature. In this paper, we focus on the evaluation of Hankel determinants featuring Catalan numbers by counting nonintersecting path systems in an associated Catalan digraph. We apply this approach in order to revisit and extend a result due to Cvetkovic-Rajkovic-Ivkovic, where we find that Hankel determinants involving the sum of successive Catalan numbers produce Fibonacci sequences.

### 1. INTRODUCTION

Combinatorial interpretations of determinants can bring deeper understanding to their evaluations; this is especially true when the entries of a matrix have natural graph theoretic descriptions. Lindström, Gessel, and Viennot [5, 6] reveal how the determinant counts signed nonintersecting path-systems in an associated directed graph. For an  $n \times n$  matrix  $A = (a_{ij})$ , the general idea is to create an acyclic directed graph  $D$  with  $n$  origin nodes,  $o_1, o_2, \dots, o_n$ , and  $n$  destination nodes,  $d_1, d_2, \dots, d_n$ , so that the number of paths from origin  $o_i$  to destination  $d_j$  is  $a_{ij}$ . Given a permutation  $\sigma$  in  $S_n$ , the product  $\prod_{i=1}^n a_{i\sigma(i)}$  counts the ways to construct  $n$  directed paths in  $D$  where the  $i^{\text{th}}$  path goes from origin  $o_i$  to destination  $d_{\sigma(i)}$ . We call such a system of  $n$  directed paths an  $n$ -route. Since  $\det(A) = \sum_{\sigma \in S_n} \text{sgn}(\sigma) \prod_{i=1}^n a_{i\sigma(i)}$ , where  $\text{sgn}(\sigma)$  is the sign of the permutation, the determinant is the number of  $n$ -routes induced by even permutations (called *even  $n$ -routes*) minus the number of  $n$ -routes induced by odd permutations (called *odd  $n$ -routes*). A sign reversing involution exists between even and odd  $n$ -routes provided some vertex of  $D$  is shared by two paths in the route, i.e. whenever two paths intersect. So calculating the determinant reduces to determining the number of even *nonintersecting  $n$ -routes* minus the number of odd *nonintersecting  $n$ -routes*.

When matrix entries are binomial coefficients, Fibonacci numbers, or a combination thereof, the nonintersecting path interpretation leads to insightful evaluations [1, 2, 10]. Catalan numbers have natural interpretations as lattice paths; consequently matrices with Catalan entries also have beautiful combinatorial explanations. Recall that the  $n^{\text{th}}$  Catalan number  $C_n = \frac{1}{n+1} \binom{2n}{n}$  counts the number of paths connecting  $(0, 0)$  to  $(n, n)$  that travel along the grid of integer lattice points of  $\mathbf{R}^2$  where each path moves up or right in one-unit steps and no path extends above the line  $y = x$  [9]. This interpretation is key to applying nonintersecting  $n$ -route

---

1991 *Mathematics Subject Classification.* 05A19, 11B39.

*Key words and phrases.* Catalan numbers, combinatorial proofs, determinants, nonintersecting paths, signed involution.

arguments to matrices containing Catalan numbers. In Section 2 we explore Mays and Wojciechowski’s work [7] calculating the determinant of the Hankel matrix  $M_n^t = (C_{i+j+t})_{i,j=0}^{n-1}$  to better familiarize the reader with the proof technique. In Section 3 we extend the ideas to calculate determinants when the matrix entries are sums of successive Catalan numbers and the determinants contain Fibonacci numbers.

## 2. HANKEL MATRICES OF CATALAN NUMBERS

Given  $n \geq 1$  and  $t \geq 0$ , define  $D_n^t$ , the *Catalan digraph with  $n$  origins and destinations at distance  $t$* , to contain the vertices of the integer lattice on and below the line  $y = x$  with arcs oriented to the right and up and origins  $o_i = (-i, -i)$  and destinations  $d_i = (i + t, i + t)$  for  $i = 0, 1, \dots, n - 1$ . See Figure 1. Notice for  $0 \leq i, j \leq n - 1$ , the number of directed paths from origin  $o_i$  to destination  $d_j$  is  $C_{t+i+j}$ .




FIGURE 1. The Catalan digraph,  $D_3^2$ . Arcs are oriented to the right and up.

Mays and Wojciechowski [7] directly argue that the determinant of the matrix

$$M_n^t = \begin{bmatrix} C_t & C_{t+1} & \cdots & C_{t+n-1} \\ C_{t+1} & C_{t+2} & \cdots & C_{t+n} \\ \vdots & & \ddots & \vdots \\ C_{t+n-1} & C_{t+n} & \cdots & C_{t+2n-2} \end{bmatrix}$$

equals the number of nonintersecting  $n$ -routes corresponding to the identity permutation on the Catalan digraph  $D_n^t$ . Because of its structure, the only nonintersecting  $n$ -routes in  $D_n^t$  correspond to the identity permutation; this leads to a clear understanding of the following determinants, which appear in [7] and we present here to motivate the new results in Section 3.

**Identity 1.** For  $n \geq 1$ ,  $\det(M_n^0) = 1$  and  $\det(M_n^1) = 1$ .

When  $o_0$  and  $d_0$  coincide ( $t = 0$ ) or their  $x$ - and  $y$ -coordinates differ by one ( $t = 1$ ), the only nonintersecting  $n$ -route in the digraph is a collection of nested right angles.

**Identity 2.** For  $n \geq 1$ ,  $\det(M_n^2) = n + 1$ .

Any nonintersecting  $n$ -route in  $D_n^2$  uses exactly  $n$  of the  $n + 1$  possible lattice points along the line  $y = 2 - x$ . Selecting which of these  $n + 1$  points to avoid uniquely determines a nonintersecting  $n$ -route.

**Identity 3.** For  $n \geq 1$ ,  $\det(M_n^3) = \sum_{k=1}^{n+1} k^2 = \frac{(n+1)(n+2)(2n+3)}{6}$ .

Any nonintersecting  $n$ -route in  $D_n^3$  uses exactly  $n$  of the  $n + 1$  possible lattice points  $\{(k + 1, -k + 2) : 1 \leq k \leq n + 1\}$  along the line  $y = 3 - x$ . For a given value of  $k$ , avoiding the point  $(k + 1, -k + 2)$  in a nonintersecting  $n$ -route requires that the directed paths from origins  $o_{k-1}, o_k, \dots, o_{n-1}$  use horizontal arcs until the line  $y = 3 - x$  and vertical arcs thereafter. The remaining  $k - 1$  paths from origins  $o_0, o_1, \dots, o_{k-2}$  use exactly  $k - 1$  of the  $k$  possible lattice points along the line  $y = 2 - x$ . The same is true for the line  $y = 4 - x$ . Selecting the points to avoid on each line, completely determines a nonintersecting  $n$ -route. See Figure 2. Since  $k$  ranges between 1 and  $n + 1$ , the determinant equals  $\sum_{k=1}^{n+1} k^2$ .




FIGURE 2. There are  $3^2$  nonintersecting 4-routes in  $D_4^3$  avoiding the point  $(4, -1)$ . A nonintersecting 4-route is determined by selecting one of three points on each line  $y = 2 - x$  and  $y = 4 - x$  to avoid.

### 3. HANKEL MATRICES OF CATALAN SUMS

The next step is to consider the determinants of Hankel matrices containing the sums of consecutive Catalan numbers. Let  $S_n^t = (C_{i+j+t} + C_{i+j+t+1})_{i,j=0}^{n-1}$ . So

$$S_n^t = \begin{bmatrix} C_t + C_{t+1} & C_{t+1} + C_{t+2} & \cdots & C_{t+n-1} + C_{t+n} \\ C_{t+1} + C_{t+2} & C_{t+2} + C_{t+3} & \cdots & C_{t+n} + C_{t+n+1} \\ \vdots & \vdots & \ddots & \vdots \\ C_{t+n-1} + C_{t+n} & C_{t+n} + C_{t+n+1} & \cdots & C_{t+2n-2} + C_{t+2n-1} \end{bmatrix}.$$

Using Hankel transforms and generating functions, Cvetković, Rajković, and Ivković [4] showed that  $\det(S_n^0) = f_{2n}$  and  $\det(S_n^1) = f_{2n+1}$  where  $f_0 = 1, f_1 = 1$ , and for  $n \geq 2, f_n = f_{n-1} + f_{n-2}$ . The simplicity of these answers begs for an elegant combinatorial solution. To see why  $\det(S_n^t)$  is a Fibonacci number for  $t = 0$  or  $1$ , we present a bijection between nonintersecting  $n$ -routes in an associated digraph and tilings of a rectangle with squares and dominoes. Recall that for  $n \geq 0$ , the Fibonacci number  $f_n$  counts the ways to tile a  $1 \times n$  rectangle using  $1 \times 1$  squares and  $1 \times 2$  dominoes [3, 8]. In fact, using this bijection with the ideas in Section 2 will allow us to take the results further and calculate  $\det(S_n^t)$  when  $t = 2$ .

We begin by defining a digraph  $\tilde{D}_n^t$  whose signed sum of nonintersecting  $n$ -routes calculates  $\det(S_n^t)$  for  $t \geq 1$ . This directed graph is obtained by adding  $n$  additional vertices,  $2n$  arcs, and relocating the destinations in the Catalan digraph  $D_n^t$ . Specifically,  $\tilde{D}_n^t$  is the digraph consisting of all the vertices of  $D_n^t$  plus the set of vertices  $\{(i+t, i+t+1) : 0 \leq i \leq n-1\}$  and the arcs of  $D_n^t$  plus vertical arcs  $\{((i+t, i+t), (i+t, i+t+1)) : \text{for } 0 \leq i \leq n-1\}$  and horizontal arcs  $\{((i+t+1, i+t+1), (i+t, i+t+1)) : \text{for } 0 \leq i \leq n-1\}$ . Notice, that the additional horizontal arcs represent steps to the left as opposed to the usual steps to the right in the Catalan digraph. Finally for  $0 \leq i \leq n-1$ , we preserve origin  $o_i = (-i, -i)$  and relocate destination  $d_i$  to the newly added vertex  $(i+t, i+t+1)$ . See Figure 3. Then the number of directed paths from origin  $o_i$  to destination  $d_j$  is  $C_{t+i+j} + C_{t+i+j+1}$  ( $C_{t+i+j}$  ways when the final step is upward and  $C_{t+i+j+1}$  ways when the final step is to the left). It is easy to see that a nonintersecting  $n$ -route in  $\tilde{D}_n^t$  can only arise from the identity permutation.




FIGURE 3. The Modified Catalan digraphs,  $\tilde{D}_3^0$  and  $\tilde{D}_3^1$ . Unmarked arcs are oriented to the right and up.

In the first identity, we consider the case when  $t = 0$  where destination  $d_0$  lies one step above origin  $o_0$ .

**Identity 4.** For  $n \geq 1, \det(S_n^0) = f_{2n}$ .

To understand this identity, we create a bijection between nonintersecting  $n$ -routes of  $\tilde{D}_n^0$  and Fibonacci tilings of a  $1 \times 2n$  board with squares and dominoes. More precisely, each nonintersecting  $n$ -route with exactly  $k$  paths taking final steps

to the left is mapped to a Fibonacci tiling of a  $1 \times 2n$  board containing exactly  $k$  dominoes.

Notice that for  $i = 0, \dots, n - 1$  a path from  $o_i$  to  $d_i$  in a nonintersecting  $n$ -route will either be “L-shaped,” taking  $2i$  steps to the right, followed by  $2i + 1$  vertical steps (and we label this path with  $\ell(i) = 0$ ) or it will take  $2i$  steps to the right, followed by  $k$  vertical steps for some  $0 \leq k \leq 2i$ . Then it completes the “hook” by taking one step right,  $2i + 1 - k$  steps up, and one final step to the left. We label this path with  $\ell(i) = 2i + 1 - k$ . Hence, when  $\ell(i) \neq 0$ ,  $\ell(i)$  is the vertical distance between the horizontal step from  $x = i$  to  $x = i + 1$  and the final left step  $((i + 1, i + 1), (i, i + 1))$ . In Figure 4,  $\ell(0) = 0$ ,  $\ell(1) = 2$ , and  $\ell(2) = 4$ . When a path in a nonintersecting  $n$ -route of  $\tilde{D}_n^0$  begins at  $o_i$  and ends with a left-step (into  $d_i$ ), the paths from origins  $o_{i+1}, o_{i+2}, \dots, o_{n-1}$  must end with left-steps as well since the vertical passage to their intended destinations are sequentially blocked. Notice that two consecutive nonzero values  $\ell(i)$  and  $\ell(i + 1)$  must differ by at least two.




FIGURE 4. The nonintersecting 3-route above with  $\ell(1) = 2$ ,  $\ell(2) = 4$  is mapped to the tiling of length 6 with dominoes beginning on cells 2 and 4.

A nonintersecting  $n$ -route will have, for some  $0 \leq k \leq n$ ,  $n - k$  L-shaped paths from  $o_i$  to  $d_i$  (for  $i = 0, \dots, n - k - 1$ ) followed by  $k$  hooked paths from  $o_i$  to  $d_i$  (for  $i = n - k, \dots, n - 1$ ). We map such an  $n$ -route to the Fibonacci tiling with  $k$  dominoes beginning on cells  $\ell(n - k), \ell(n - k + 1), \dots, \ell(n - 1)$  and squares everywhere else. Since  $0 \leq \ell(n - 1) \leq 2n - 1$ , a final domino can be begin on cell  $2n - 1$  and the Fibonacci tiling has length  $2n$ .

Every nonintersecting  $n$ -route is mapped to a unique tiling of the  $1 \times 2n$  rectangle. Conversely, every tiling of a  $1 \times 2n$  rectangle (with  $k$  dominoes) induces a unique nonintersecting  $n$ -route since the position of the dominoes defines the locations of the final steps to the right for (the last  $k$ ) paths ending with left-steps. Thus, a bijection exists and  $\det(S_n^0)$  equals the number of Fibonacci tilings of length  $2n$ , namely  $f_{2n}$ .

**Identity 5.** For  $n \geq 1$ ,  $\det(S_n^1) = f_{2n+1}$ .

The bijection is essentially the same as above. A nonintersecting  $n$ -route in  $\tilde{D}_n^1$  whose  $k$  paths from origins  $o_{n-k}, o_{n-k+1}, \dots, o_{n-1}$  utilize the final left-steps is



- [2] A. T. Benjamin, N. T. Cameron, J.J. Quinn, Fibonacci determinants—a combinatorial approach, *Fibonacci Quarterly*, to appear.
- [3] A. T. Benjamin and J. J. Quinn, *Proofs That Really Count: The Art of Combinatorial Proof*, Mathematical Association of America, Washington, D.C., 2003.
- [4] A. Cvetković, P. Rajković, M. Ivković, Catalan numbers, the Hankel transform, and Fibonacci numbers, *Journal of Integer Sequences*, Vol. 5 (2002), Article 02.1.3.
- [5] I. Gessel and G. Viennot, Binomial determinants, paths, and hook length formulae. *Adv. in Math.* **58** (1985), no. 3, 300–321.
- [6] B. Lindström, On the vector representations of induced matroids, *Bull. London Math. Soc.* **5** (1973) 85–90.
- [7] M.E. Mays and J. Wojciechowski, A determinant property of Catalan numbers. *Discrete Math.* **211** (2000), no. 1-3, 125–133.
- [8] R. Stanley, *Enumerative Combinatorics, Volume 1*, Wadsworth & Brooks/Cole, California, 1986.
- [9] R. Stanley, *Enumerative Combinatorics, Volume 2*, Cambridge University Press, 1997.
- [10] D. Zeilberger, A combinatorial approach to matrix algebra, *Discrete Math.* **56** (1985) 61–72.

HARVEY MUDD COLLEGE  
*E-mail address:* [benjamin@hmc.edu](mailto:benjamin@hmc.edu)

LEWIS & CLARK COLLEGE  
*E-mail address:* [ncameron@lclark.edu](mailto:ncameron@lclark.edu)

ASSOCIATION FOR WOMEN IN MATHEMATICS  
*E-mail address:* [jquinn@awm-math.org](mailto:jquinn@awm-math.org)

HARVEY MUDD COLLEGE  
*E-mail address:* [carl\\_yerger@math.hmc.edu](mailto:carl_yerger@math.hmc.edu)