

See all >
[13 Citations](#)

See all >
[34 References](#)


[Share](#)

[Download Full-text PDF](#)

On a class of q -Bernoulli and q -Euler polynomials

Article (PDF Available) in *Advances in Difference Equations* 2013(1) · April 2013 with 36 Reads

Impact Factor: 0.64 · DOI: 10.1186/1687-1847-2013-108



1st [Nazim Mahmudov](#)
32.56 · Eastern Mediterranean Un...




Image Analyser G S A

A program for scientific evaluation of 2D images

- ✓ Object surface calculation
- ✓ Object length calculation
- ✓ Object counting
- ✓ And much more

[Download Free Trial](#)

Abstract

The main purpose of this paper is to introduce and investigate a class of generalized Bernoulli polynomials and Euler polynomials based on the q -integers. The q -analogues of well-known formulas are derived. The q -analogue of the Srivastava-Pintér addition theorem is obtained. We give new identities involving q -Bernstein polynomials.

Discover the world's research

- **100 million** publications
- **2.5 million** new publications each month
- **10 million** members

[Join for free](#)

Full-text (PDF)

Available from: [Nazim Mahmudov](#), May 07, 2014

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · *Advances in Difference Equations*

RESEARCH

Open Access

On a class of q -Bernoulli and q -Euler polynomials

Nazim I Mahmudov*

*Correspondence:
nazim.mahmudov@emu.edu.tr
Eastern Mediterranean University,
Gazimagusa, T.R. North Cyprus, via
Mersin 10, Turkey

Abstract

The main purpose of this paper is to introduce and investigate a class of generalized Bernoulli polynomials and Euler polynomials based on the q -integers. The q -analogues of well-known formulas are derived. The q -analogue of the Srivastava-Pintér addition theorem is obtained. We give new identities involving q -Bernstein polynomials.

1 Introduction

Throughout this paper, we always make use of the following notation: \mathbb{N} denotes the set of natural numbers, \mathbb{N}_0 denotes the set of nonnegative integers, \mathbb{R} denotes the set of real numbers, \mathbb{C} denotes the set of complex numbers.

The q -shifted factorial is defined by

$$(a; q)_0 = 1, \quad (a; q)_n = \prod_{j=0}^{n-1} (1 - q^j a), \quad n \in \mathbb{N},$$

$$(a; q)_\infty = \prod_{j=0}^{\infty} (1 - q^j a), \quad |q| < 1, a \in \mathbb{C}.$$

The q -numbers and q -numbers factorial are defined by

$$[a]_q = \frac{1 - q^a}{1 - q} \quad (q \neq 1); \quad [0]_q! = 1; \quad [n]_q! = [1]_q [2]_q \cdots [n]_q, \quad n \in \mathbb{N}, a \in \mathbb{C},$$

respectively. The q -polynomial coefficient is defined by

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{(q; q)_n}{(q; q)_{n-k} (q; q)_k}.$$

The q -analogue of the function $(x + y)^n$ is defined by

$$(x + y)_q^n := \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{\frac{1}{2}k(k-1)} x^{n-k} y^k, \quad n \in \mathbb{N}_0.$$

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equ...

The q -binomial formula is known as

$$(1-a)_q^n = (a; q)_n = \prod_{j=0}^{n-1} (1-q^j a) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{\frac{1}{2}k(k-1)} (-1)^k a^k.$$

In the standard approach to the q -calculus, two exponential functions are used

$$e_q(z) = \sum_{n=0}^{\infty} \frac{z^n}{[n]_q!} = \prod_{k=0}^{\infty} \frac{1}{(1-(1-q)q^k z)}, \quad 0 < |q| < 1, |z| < \frac{1}{|1-q|},$$

$$E_q(z) = \sum_{n=0}^{\infty} \frac{q^{\frac{1}{2}n(n-1)} z^n}{[n]_q!} = \prod_{k=0}^{\infty} (1 + (1-q)q^k z), \quad 0 < |q| < 1, z \in \mathbb{C}.$$

From this form, we easily see that $e_q(z)E_q(-z) = 1$. Moreover,

$$D_q e_q(z) = e_q(z), \quad D_q E_q(z) = E_q(qz),$$

where D_q is defined by

$$D_q f(z) := \frac{f(qz) - f(z)}{qz - z}, \quad 0 < |q| < 1, 0 \neq z \in \mathbb{C}.$$

The above q -standard notation can be found in [1].

Over 70 years ago, Carlitz extended the classical Bernoulli and Euler numbers and polynomials and introduced the q -Bernoulli and the q -Euler numbers and polynomials (see [2, 3] and [4]). There are numerous recent investigations on this subject by, among many other authors, Cenkci *et al.* [5–7], Choi *et al.* [8] and [9], Kim *et al.* [10–13], Ozden and Simsek [14], Ryoo *et al.* [15], Simsek [16, 17] and [18], and Luo and Srivastava [19], Srivastava *et al.* [20], Srivastava [21], Mahmudov [22].

We first give here the definitions of the q -Bernoulli and the q -Euler polynomials of higher order as follows.

Definition 1 Let $q, \alpha \in \mathbb{C}$, $0 < |q| < 1$. The q -Bernoulli numbers $\mathfrak{B}_{n,q}^{(\alpha)}$ and polynomials $\mathfrak{B}_{n,q}^{(\alpha)}(x, y)$ in x, y of order α are defined by means of the generating functions:

$$\left(\frac{t}{e_q(t) - 1} \right)^\alpha = \sum_{n=0}^{\infty} \mathfrak{B}_{n,q}^{(\alpha)} \frac{t^n}{[n]_q!}, \quad |t| < 2\pi,$$

$$\left(\frac{t}{e_q(t) - 1} \right)^\alpha e_q(tx) E_q(ty) = \sum_{n=0}^{\infty} \mathfrak{B}_{n,q}^{(\alpha)}(x, y) \frac{t^n}{[n]_q!}, \quad |t| < 2\pi.$$

Definition 2 Let $q, \alpha \in \mathbb{C}$, $0 < |q| < 1$. The q -Euler numbers $\mathfrak{E}_{n,q}^{(\alpha)}$ and polynomials $\mathfrak{E}_{n,q}^{(\alpha)}(x, y)$ in x, y of order α are defined by means of the generating functions:

$$\left(\frac{2}{e_q(t) + 1} \right)^\alpha = \sum_{n=0}^{\infty} \mathfrak{E}_{n,q}^{(\alpha)} \frac{t^n}{[n]_q!}, \quad |t| < \pi,$$

$$\left(\frac{2}{e_q(t) + 1} \right)^\alpha e_q(t) \frac{t^n}{[n]_q!}$$

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equ...

People who read this publication also read:

Article: [On a class of generalized \$q\$ -Bernoulli and \$q\$ -Euler polynomials](#)

Full-text · Apr 2013 · Advances in Difference Equ...

It is obvious that

$$\begin{aligned} \mathfrak{B}_{n,q}^{(\alpha)} &= \mathfrak{B}_{n,q}^{(\alpha)}(0, 0), & \lim_{q \rightarrow 1^-} \mathfrak{B}_{n,q}^{(\alpha)}(x, y) &= B_n^{(\alpha)}(x + y), & \lim_{q \rightarrow 1^-} \mathfrak{B}_{n,q}^{(\alpha)} &= B_n^{(\alpha)}, \\ \mathfrak{E}_{n,q}^{(\alpha)} &= \mathfrak{E}_{n,q}^{(\alpha)}(0, 0), & \lim_{q \rightarrow 1^-} \mathfrak{E}_{n,q}^{(\alpha)}(x, y) &= E_n^{(\alpha)}(x + y), & \lim_{q \rightarrow 1^-} \mathfrak{E}_{n,q}^{(\alpha)} &= E_n^{(\alpha)}, \\ \lim_{q \rightarrow 1^-} \mathfrak{B}_{n,q}^{(\alpha)}(x, 0) &= B_n^{(\alpha)}(x), & \lim_{q \rightarrow 1^-} \mathfrak{B}_{n,q}^{(\alpha)}(0, y) &= B_n^{(\alpha)}(y), \\ \lim_{q \rightarrow 1^-} \mathfrak{E}_{n,q}^{(\alpha)}(x, 0) &= E_n^{(\alpha)}(x), & \lim_{q \rightarrow 1^-} \mathfrak{E}_{n,q}^{(\alpha)}(0, y) &= E_n^{(\alpha)}(y). \end{aligned}$$

Here $B_n^{(\alpha)}(x)$ and $E_n^{(\alpha)}(x)$ denote the classical Bernoulli and Euler polynomials of order α which are defined by

$$\left(\frac{t}{e^t - 1}\right)^\alpha e^{tx} = \sum_{n=0}^{\infty} B_n^{(\alpha)}(x) \frac{t^n}{n!} \quad \text{and} \quad \left(\frac{2}{e^t + 1}\right)^\alpha e^{tx} = \sum_{n=0}^{\infty} E_n^{(\alpha)}(x) \frac{t^n}{n!}.$$

In fact, Definitions 1 and 2 define two different types $\mathfrak{B}_{n,q}^{(\alpha)}(x, 0)$ and $\mathfrak{B}_{n,q}^{(\alpha)}(0, y)$ of the q -Bernoulli polynomials and two different types $\mathfrak{E}_{n,q}^{(\alpha)}(x, 0)$ and $\mathfrak{E}_{n,q}^{(\alpha)}(0, y)$ of the q -Euler polynomials. Both polynomials $\mathfrak{B}_{n,q}^{(\alpha)}(x, 0)$ and $\mathfrak{B}_{n,q}^{(\alpha)}(0, y)$ ($\mathfrak{E}_{n,q}^{(\alpha)}(x, 0)$ and $\mathfrak{E}_{n,q}^{(\alpha)}(0, y)$) coincide with the classical higher-order Bernoulli polynomials (Euler polynomials) in the limiting case $q \rightarrow 1^-$.

For the q -Bernoulli numbers $\mathfrak{B}_{n,q}$ and the q -Euler numbers $\mathfrak{E}_{n,q}$ of order n , we have

$$\mathfrak{B}_{n,q} = \mathfrak{B}_{n,q}(0, 0) = \mathfrak{B}_{n,q}^{(1)}(0, 0), \quad \mathfrak{E}_{n,q} = \mathfrak{E}_{n,q}(0, 0) = \mathfrak{E}_{n,q}^{(1)}(0, 0),$$

respectively. Note that the q -Bernoulli numbers $\mathfrak{B}_{n,q}$ are defined and studied in [23].

The aim of the present paper is to obtain some results for the above newly defined q -Bernoulli and q -Euler polynomials. It should be mentioned that q -Bernoulli and q -Euler polynomials in our definitions are polynomials of x and y , and when $y = 0$, they are polynomials of x , but in other definitions they are functions of q^x . First advantage of this approach is that for $q \rightarrow 1^-$, $\mathfrak{B}_{n,q}^{(\alpha)}(x, y)$ ($\mathfrak{E}_{n,q}^{(\alpha)}(x, y)$) becomes the classical Bernoulli $B_n^{(\alpha)}(x + y)$ (Euler $E_n^{(\alpha)}(x + y)$) polynomial, and we may obtain the q -analogues of well-known results, for example, those of Srivastava and Pintér [24], Cheon [25], etc. The second advantage is that we find the relation between q -Bernstein polynomials and Phillips q -Bernoulli polynomials and derive the formulas involving the q -Stirling numbers of the second kind, q -Bernoulli polynomials and Phillips q -Bernstein polynomials.

2 Preliminaries and lemmas

In this section we provide some basic formulas for the q -Bernoulli and q -Euler polynomials in order to obtain the main results of this paper in the next section. The following result is a q -analogue of the addition theorem for the classical Bernoulli and Euler polynomials.

Lemma 3 (Addition theorems) *For all $x, y \in \mathbb{C}$, we have*

$$\mathfrak{B}_{n,q}^{(\alpha)}(x, y) = \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_q \mathfrak{B}_{k,q}^{(\alpha)}(x, y) \mathfrak{B}_{n-k,q}^{(\alpha)}(0, y)$$

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equations

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equ...

$$\mathfrak{B}_{n,q}^{(\alpha)}(x, y) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{(n-k)(n-k-1)/2} \mathfrak{B}_{k,q}^{(\alpha)}(x, 0) y^{n-k} = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathfrak{B}_{k,q}^{(\alpha)}(0, y) x^{n-k}, \quad (1)$$

$$\mathfrak{E}_{n,q}^{(\alpha)}(x, y) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{(n-k)(n-k-1)/2} \mathfrak{E}_{k,q}^{(\alpha)}(x, 0) y^{n-k} = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathfrak{E}_{k,q}^{(\alpha)}(0, y) x^{n-k}. \quad (2)$$

In particular, setting $x = 0$ and $y = 0$ in (1) and (2), we get the following formulas for q -Bernoulli and q -Euler polynomials, respectively

$$\mathfrak{B}_{n,q}^{(\alpha)}(x, 0) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathfrak{B}_{k,q}^{(\alpha)} x^{n-k}, \quad \mathfrak{B}_{n,q}^{(\alpha)}(0, y) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{(n-k)(n-k-1)/2} \mathfrak{B}_{k,q}^{(\alpha)} y^{n-k}, \quad (3)$$

$$\mathfrak{E}_{n,q}^{(\alpha)}(x, 0) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathfrak{E}_{k,q}^{(\alpha)} x^{n-k}, \quad \mathfrak{E}_{n,q}^{(\alpha)}(0, y) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{(n-k)(n-k-1)/2} \mathfrak{E}_{k,q}^{(\alpha)} y^{n-k}. \quad (4)$$

Setting $y = 1$ and $x = 1$ in (1) and (2), we get

$$\mathfrak{B}_{n,q}^{(\alpha)}(x, 1) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{(n-k)(n-k-1)/2} \mathfrak{B}_{k,q}^{(\alpha)}(x, 0), \quad \mathfrak{B}_{n,q}^{(\alpha)}(1, y) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathfrak{B}_{k,q}^{(\alpha)}(0, y), \quad (5)$$

$$\mathfrak{E}_{n,q}^{(\alpha)}(x, 1) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{(n-k)(n-k-1)/2} \mathfrak{E}_{k,q}^{(\alpha)}(x, 0), \quad \mathfrak{E}_{n,q}^{(\alpha)}(1, y) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathfrak{E}_{k,q}^{(\alpha)}(0, y). \quad (6)$$

Clearly, (5) and (6) are q -analogues of

$$B_n^{(\alpha)}(x+1) = \sum_{k=0}^n \binom{n}{k} B_k^{(\alpha)}(x), \quad E_n^{(\alpha)}(x+1) = \sum_{k=0}^n \binom{n}{k} E_k^{(\alpha)}(x),$$

respectively.

Lemma 4 *We have*

$$D_{q,x} \mathfrak{B}_{n,q}^{(\alpha)}(x, y) = [n]_q \mathfrak{B}_{n-1,q}^{(\alpha)}(x, y), \quad D_{q,y} \mathfrak{B}_{n,q}^{(\alpha)}(x, y) = [n]_q \mathfrak{B}_{n-1,q}^{(\alpha)}(x, qy),$$

$$D_{q,x} \mathfrak{E}_{n,q}^{(\alpha)}(x, y) = [n]_q \mathfrak{E}_{n-1,q}^{(\alpha)}(x, y), \quad D_{q,y} \mathfrak{E}_{n,q}^{(\alpha)}(x, y) = [n]_q \mathfrak{E}_{n-1,q}^{(\alpha)}(x, qy).$$

Lemma 5 (Difference equations) *We have*

$$\mathfrak{B}_{n,q}^{(\alpha)}(1, y) - \mathfrak{B}_{n,q}^{(\alpha)}(0, y) = [n]_q \mathfrak{B}_{n-1,q}^{(\alpha-1)}(0, y), \quad (7)$$

$$\mathfrak{E}_{n,q}^{(\alpha)}(1, y) + \mathfrak{E}_{n,q}^{(\alpha)}(0, y) = 2 \mathfrak{E}_{n,q}^{(\alpha-1)}(0, y), \quad (8)$$

$$\mathfrak{B}_{n,q}^{(\alpha)}(x, 0) - \mathfrak{B}_{n,q}^{(\alpha)}(x, -1) = [n]_q \mathfrak{B}_{n-1,q}^{(\alpha-1)}(x, -1),$$

$$\mathfrak{E}_{n,q}^{(\alpha)}(x, 0) + \mathfrak{E}_{n,q}^{(\alpha)}(x, -1) = 2 \mathfrak{E}_{n,q}^{(\alpha-1)}(x, -1).$$

From (7) and (5), (8) and (6), we obtain the following formulas.

Lemma 6 *We have*

$$\mathfrak{B}_{n,q}^{(\alpha-1)}(0, y) =$$

People who read this publication also read:

[Article: On a class of generalized \$q\$ -Bernoulli and \$q\$ -Euler polynomials](#)

Full-text · Apr 2013 · *Advances in Difference Equations*

People who read this publication also read:

Article: [On a class of generalized \$q\$ -Bernoulli and \$q\$ -Euler polynomials](#)

Full-text · Apr 2013 · Advances in Difference Equ...

$$\mathfrak{E}_{n,q}^{(\alpha-1)}(0,y) = \frac{1}{2} \left[\sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathfrak{E}_{k,q}^{(\alpha)}(0,y) + \mathfrak{E}_{n,q}^{(\alpha)}(0,y) \right]. \quad (10)$$

Putting $\alpha = 1$ in (9) and (10) and noting that

$$\mathfrak{B}_{n,q}^{(0)}(0,y) = \mathfrak{E}_{n,q}^{(0)}(0,y) = q^{n(n-1)/2} y^n,$$

we arrive at the following expansions:

$$y^n = \frac{1}{q^{n(n-1)/2} [n+1]_q} \sum_{k=0}^n \begin{bmatrix} n+1 \\ k \end{bmatrix}_q \mathfrak{B}_{k,q}(0,y),$$

$$y^n = \frac{1}{2q^{n(n-1)/2}} \left[\sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathfrak{E}_{k,q}(0,y) + \mathfrak{E}_{n,q}(0,y) \right],$$

which are q -analogues of the following familiar expansions:

$$y^n = \frac{1}{n+1} \sum_{k=0}^n \binom{n+1}{k} B_k(y), \quad y^n = \frac{1}{2} \left[\sum_{k=0}^n \binom{n}{k} E_k(y) + E_n(y) \right], \quad (11)$$

respectively.

Lemma 7 (Recurrence relationships) *The polynomials $\mathfrak{B}_{n,q}^{(\alpha)}(x,0)$ and $\mathfrak{E}_{n,q}^{(\alpha)}(x,0)$ satisfy the following difference relationships:*

$$\sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix}_q m^j \mathfrak{B}_{j,q}^{(\alpha)}(x,0) - \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix}_q m^j \mathfrak{B}_{j,q}^{(\alpha)}(x,-1)$$

$$= [k]_q \sum_{j=0}^{k-1} \begin{bmatrix} k-1 \\ j \end{bmatrix}_q m^{j+1} \mathfrak{B}_{j,q}^{(\alpha-1)}(x,-1), \quad (12)$$

$$\mathfrak{B}_{k,q}^{(\alpha)}\left(\frac{1}{m}, y\right) - \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix}_q \left(\frac{1}{m} - 1\right)_q^{k-j} \mathfrak{B}_{j,q}^{(\alpha)}(0,y)$$

$$= [k]_q \sum_{j=0}^{k-1} \begin{bmatrix} k-1 \\ j \end{bmatrix}_q \left(\frac{1}{m} - 1\right)_q^{k-j-1} \mathfrak{B}_{j,q}^{(\alpha-1)}(0,y), \quad (13)$$

$$\sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix}_q m^j \mathfrak{E}_{j,q}^{(\alpha)}(x,0) + \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix}_q m^j \mathfrak{E}_{j,q}^{(\alpha)}(x,-1)$$

$$= 2 \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix}_q m^j \mathfrak{E}_{j,q}^{(\alpha-1)}(x,-1), \quad (14)$$

$$\mathfrak{E}_{k,q}^{(\alpha)}\left(\frac{1}{m}, y\right) + \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix}_q \left(\frac{1}{m} - 1\right)_q^{k-j} \mathfrak{E}_{j,q}^{(\alpha)}(0,y)$$

$$= 2 \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix}_q \left(\frac{1}{m} - 1\right)_q^{k-j} \mathfrak{E}_{j,q}^{(\alpha-1)}(0,y)$$

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equations

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equ...

3 Explicit relationship between the q -Bernoulli and q -Euler polynomials

In this section we investigate some explicit relationships between the q -Bernoulli and q -Euler polynomials. Here some q -analogues of known results are given. We also obtain new formulas and some of their special cases below. These formulas are some extensions of the formulas of Srivastava and Pintér, Cheon and others.

We present natural q -extensions of the main results in the papers [24] and [26], see Theorems 8 and 13.

Theorem 8 For $n \in \mathbb{N}_0$, the following relationships hold true:

$$\begin{aligned} \mathfrak{B}_{n,q}^{(\alpha)}(x,y) &= \frac{1}{2m^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \left[m^k \mathfrak{B}_{k,q}^{(\alpha)}(x,0) + \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix}_q m^j \mathfrak{B}_{j,q}^{(\alpha)}(x,-1) \right. \\ &\quad \left. + [k]_q \sum_{j=0}^{k-1} \begin{bmatrix} k-1 \\ j \end{bmatrix}_q m^{j+1} \mathfrak{B}_{j,q}^{(\alpha-1)}(x,-1) \right] \mathfrak{E}_{n-k,q}(0,my), \\ \mathfrak{B}_{n,q}^{(\alpha)}(x,y) &= \frac{1}{2m^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q m^k \left[\mathfrak{B}_{k,q}^{(\alpha)}(0,y) + \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix}_q \left(\frac{1}{m} - 1 \right)_q^{k-j} \mathfrak{B}_{j,q}^{(\alpha)}(0,y) \right. \\ &\quad \left. + [k]_q \sum_{j=0}^{k-1} \begin{bmatrix} k-1 \\ j \end{bmatrix}_q \left(\frac{1}{m} - 1 \right)_q^{k-1-j} \mathfrak{B}_{j,q}^{(\alpha-1)}(0,y) \right] \mathfrak{E}_{n-k,q}(mx,0). \end{aligned}$$

Proof Using the following identity:

$$\begin{aligned} &\left(\frac{t}{e_q(t)-1} \right)^\alpha e_q(tx) E_q(ty) \\ &= \frac{2}{e_q\left(\frac{t}{m}\right)+1} \cdot E_q\left(\frac{t}{m} my\right) \cdot \frac{e_q\left(\frac{t}{m}\right)+1}{2} \cdot \left(\frac{t}{e_q(t)-1} \right)^\alpha e_q(tx), \end{aligned}$$

we have

$$\begin{aligned} \sum_{n=0}^{\infty} \mathfrak{B}_{n,q}^{(\alpha)}(x,y) \frac{t^n}{[n]_q!} &= \frac{1}{2} \sum_{n=0}^{\infty} \mathfrak{E}_{n,q}(0,my) \frac{t^n}{m^n [n]_q!} \sum_{n=0}^{\infty} \frac{t^n}{m^n [n]_q!} \sum_{n=0}^{\infty} \mathfrak{B}_{n,q}^{(\alpha)}(x,0) \frac{t^n}{[n]_q!} \\ &\quad + \frac{1}{2} \sum_{n=0}^{\infty} \mathfrak{E}_{n,q}(0,my) \frac{t^n}{m^n [n]_q!} \sum_{n=0}^{\infty} \mathfrak{B}_{n,q}^{(\alpha)}(x,0) \frac{t^n}{[n]_q!} \\ &=: I_1 + I_2. \end{aligned}$$

It is clear that

$$\begin{aligned} I_2 &= \frac{1}{2} \sum_{n=0}^{\infty} \mathfrak{E}_{n,q}(0,my) \frac{t^n}{m^n [n]_q!} \sum_{n=0}^{\infty} \mathfrak{B}_{n,q}^{(\alpha)}(x,0) \frac{t^n}{[n]_q!} \\ &= \frac{1}{2} \sum_{n=0}^{\infty} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \end{aligned}$$

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equ...

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equ...

On the other hand,

$$\begin{aligned}
 I_1 &= \frac{1}{2} \sum_{n=0}^{\infty} \mathfrak{B}_{n,q}^{(\alpha)}(x, 0) \frac{t^n}{[n]_q!} \sum_{n=0}^{\infty} \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}_q m^{-n} \mathfrak{E}_{j,q}(0, my) \frac{t^n}{[n]_q!} \\
 &= \frac{1}{2} \sum_{n=0}^{\infty} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathfrak{B}_{k,q}^{(\alpha)}(x, 0) \sum_{j=0}^{n-k} \begin{bmatrix} n-k \\ j \end{bmatrix}_q m^{k-n} \mathfrak{E}_{j,q}(0, my) \frac{t^n}{[n]_q!} \\
 &= \frac{1}{2} \sum_{n=0}^{\infty} m^{-n} \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}_q \mathfrak{E}_{j,q}(0, my) \sum_{k=0}^{n-j} \begin{bmatrix} n-j \\ k \end{bmatrix}_q m^k \mathfrak{B}_{k,q}^{(\alpha)}(x, 0) \frac{t^n}{[n]_q!}.
 \end{aligned}$$

Therefore

$$\begin{aligned}
 &\sum_{n=0}^{\infty} \mathfrak{B}_{n,q}^{(\alpha)}(x, y) \frac{t^n}{[n]_q!} \\
 &= \frac{1}{2} \sum_{n=0}^{\infty} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q m^{k-n} \left[\mathfrak{B}_{k,q}^{(\alpha)}(x, 0) + m^{-k} \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix}_q m^j \mathfrak{B}_{j,q}^{(\alpha)}(x, 0) \right] \mathfrak{E}_{n-k,q}(0, my) \frac{t^n}{[n]_q!}.
 \end{aligned}$$

It remains to use the formula (12). □

Next we discuss some special cases of Theorem 8.

Corollary 9 For $n \in \mathbb{N}_0$, $m \in \mathbb{N}$, the relationship

$$\begin{aligned}
 \mathfrak{B}_{n,q}(x, y) &= \frac{1}{2m^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \left[m^k \mathfrak{B}_{k,q}(x, 0) + \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix}_q m^j \mathfrak{B}_{j,q}(x, -1) \right. \\
 &\quad \left. + [k]_q \sum_{j=0}^{k-1} \begin{bmatrix} k-1 \\ j \end{bmatrix}_q m^{j+1} (x-1)_q^j \right] \mathfrak{E}_{n-k,q}(0, my), \\
 \mathfrak{B}_{n,q}(x, y) &= \frac{1}{2m^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \left[m^k \mathfrak{B}_{k,q}(0, y) + \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix}_q \left(\frac{1}{m} - 1 \right)_q^{k-j} \mathfrak{B}_{j,q}(0, y) \right. \\
 &\quad \left. + [k]_q \sum_{j=0}^{k-1} \begin{bmatrix} k-1 \\ j \end{bmatrix}_q q^{\frac{1}{2}j(j-1)} \left(\frac{1}{m} - 1 \right)_q^{k-1-j} y^j \right] \mathfrak{E}_{n-k,q}(mx, 0)
 \end{aligned}$$

holds true between the q -Bernoulli polynomials and q -Euler polynomials.

Corollary 10 ([26]) For $n \in \mathbb{N}_0$, $m \in \mathbb{N}$, the following relationship holds true:

$$\begin{aligned}
 B_n(x+y) &= \sum_{k=0}^n \binom{n}{k} \left(B_k(y) + \frac{k}{2} y^{k-1} \right) E_{n-k}(x), \\
 B_n(x+y) &= \frac{1}{2m^n} \sum_{k=0}^n \binom{n}{k} \left[m^k B_k(x) + m^k B_k \left(x - 1 + \frac{1}{m} \right) \right.
 \end{aligned}$$

+ km

People who read this publication also read:

[Article: On a class of generalized \$q\$ -Bernoulli and \$q\$ -Euler polynomials](#)

Full-text · Apr 2013 · Advances in Difference Equ...

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equ...

Corollary 11 For $n \in \mathbb{N}_0$, the following relationship holds true:

$$\mathfrak{B}_{n,q}(x, y) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \left(\mathfrak{B}_{k,q}(0, y) + q^{\frac{1}{2}(k-1)(k-2)} \begin{bmatrix} k \\ 2 \end{bmatrix}_q y^{k-1} \right) \mathfrak{E}_{n-k,q}(x, 0). \quad (16)$$

Corollary 12 For $n \in \mathbb{N}_0$, the following relationship holds true:

$$\mathfrak{B}_{n,q}(x, 0) = \sum_{\substack{k=0 \\ (k \neq 1)}}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathfrak{B}_{k,q} \mathfrak{E}_{n-k,q}(x, 0) + \left(\mathfrak{B}_{1,q} + \frac{1}{2} \right) \mathfrak{E}_{n-1,q}(x, 0), \quad (17)$$

$$\mathfrak{B}_{n,q}(0, y) = \sum_{\substack{k=0 \\ (k \neq 1)}}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathfrak{B}_{k,q} \mathfrak{E}_{n-k,q}(0, y) + \left(\mathfrak{B}_{1,q} + \frac{1}{2} \right) \mathfrak{E}_{n-1,q}(0, y). \quad (18)$$

The formulas (16)-(18) are the q -extension of Cheon's main result [25]. Notice that $\mathfrak{B}_{1,q} = -\frac{1}{[2]_q}$, see [23], and the extra term becomes zero for $q \rightarrow 1^-$.

Theorem 13 For $n \in \mathbb{N}_0$, the relationships

$$\begin{aligned} \mathfrak{E}_{n,q}^{(\alpha)}(x, y) &= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q m^{n-1} [k+1]_q \left[2 \sum_{j=0}^{k+1} \begin{bmatrix} k+1 \\ j \end{bmatrix}_q \left(\frac{1}{m} - 1 \right)_q^{k+1-j} \mathfrak{E}_{j,q}^{(\alpha-1)}(0, y) \right. \\ &\quad \left. - \sum_{j=0}^{k+1} \begin{bmatrix} k+1 \\ j \end{bmatrix}_q \left(\frac{1}{m} - 1 \right)_q^{k+1-j} \mathfrak{E}_{j,q}^{(\alpha)}(0, y) - \mathfrak{E}_{k+1,q}^{(\alpha)}(0, y) \right] \mathfrak{B}_{n-k,q}(mx, 0), \\ \mathfrak{E}_{n,q}^{(\alpha)}(x, y) &= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q m^n [k+1]_q \left[2 \sum_{j=0}^{k+1} \begin{bmatrix} k+1 \\ j \end{bmatrix}_q m^j \mathfrak{E}_{j,q}^{(\alpha-1)}(x, -1) \right. \\ &\quad \left. - \sum_{j=0}^{k+1} \begin{bmatrix} k+1 \\ j \end{bmatrix}_q m^j \mathfrak{E}_{j,q}^{(\alpha)}(x, -1) - m^{k+1} \mathfrak{E}_{k+1,q}^{(\alpha)}(x, 0) \right] \mathfrak{B}_{n-k,q}(0, my) \end{aligned}$$

hold true between the q -Bernoulli polynomials and q -Euler polynomials.

Proof The proof is based on the following identities:

$$\begin{aligned} \left(\frac{2}{e_q(t) + 1} \right)^\alpha e_q(tx) E_q(ty) &= \left(\frac{2}{e_q(t) + 1} \right)^\alpha E_q(ty) \cdot \frac{e_q\left(\frac{t}{m}\right) - 1}{t} \cdot \frac{t}{e_q\left(\frac{t}{m}\right) - 1} e_q\left(\frac{t}{m} mx\right), \\ \left(\frac{2}{e_q(t) + 1} \right)^\alpha e_q(tx) E_q(ty) &= \left(\frac{2}{e_q(t) + 1} \right)^\alpha e_q(tx) \cdot \frac{e_q\left(\frac{t}{m}\right) - 1}{t} \cdot \frac{t}{e_q\left(\frac{t}{m}\right) - 1} E_q\left(\frac{t}{m} my\right) \end{aligned}$$

and is similar to that of Theorem 8. □

Next we discuss sor

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equ...

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equ...

Corollary 14 For $n \in \mathbb{N}_0$, $m \in \mathbb{N}$, the relationship

$$\begin{aligned} \mathfrak{E}_{n,q}(x, y) = & \sum_{k=0}^n \binom{n}{k}_q m^{-n} \left[\begin{matrix} k+1 \\ j \end{matrix} \right]_q m^j (x-1)_q^j \\ & - \sum_{j=0}^{k+1} \left[\begin{matrix} k+1 \\ j \end{matrix} \right]_q m^j \mathfrak{E}_{j,q}(x, -1) - m^{k+1} \mathfrak{E}_{k+1,q}(x, 0) \Big] \mathfrak{B}_{n-k,q}(0, my) \end{aligned}$$

holds true between the q -Bernoulli polynomials and q -Euler polynomials.

Corollary 15 ([26]) For $n \in \mathbb{N}_0$, $m \in \mathbb{N}$, the following relationships hold true:

$$\begin{aligned} E_n(x+y) &= \sum_{k=0}^n \frac{2}{k+1} \binom{n}{k} (y^{k+1} - E_{k+1}(y)) B_{n-k}(x), \\ E_n(x+y) &= \sum_{k=0}^n \binom{n}{k} \frac{m^{k-n+1}}{k+1} \left[2 \binom{x+1-m}{m}^{k+1} - E_{k+1} \left(\frac{x+1-m}{m} \right) \right. \\ & \quad \left. - E_{k+1}(x) \right] B_{n-k}(my). \end{aligned}$$

Corollary 16 For $n \in \mathbb{N}_0$, the following relationship holds true:

$$\mathfrak{E}_{n,q}(x, y) = \sum_{k=0}^n \binom{n}{k}_q \frac{2}{[k+1]_q} (q^{2^{k(k+1)}} y^{k+1} - \mathfrak{E}_{k+1,q}(0, y)) \mathfrak{B}_{n-k,q}(x, 0).$$

Corollary 17 For $n \in \mathbb{N}_0$, the following relationships hold true:

$$\begin{aligned} \mathfrak{E}_{n,q}(x, 0) &= - \sum_{k=0}^n \binom{n}{k}_q \frac{2}{[k+1]_q} \mathfrak{E}_{k+1,q} \mathfrak{B}_{n-k,q}(x, 0), \\ \mathfrak{E}_{n,q}(0, y) &= - \sum_{k=0}^n \binom{n}{k}_q \frac{2}{[k+1]_q} \mathfrak{E}_{k+1,q} \mathfrak{B}_{n-k,q}(0, y). \end{aligned}$$

These formulas are q -analogues of the formula of Srivastava and Pintér [24].

4 q -Stirling numbers and q -Bernoulli polynomials

In this section, we aim to derive several formulas involving the q -Bernoulli polynomials, the q -Euler polynomials of order α , the q -Stirling numbers of the second kind and the q -Bernstein polynomials.

Theorem 18 Each of the following relationships holds true for the Stirling numbers $S_2(n, k)$ of the second kind:

$$\begin{aligned} \mathfrak{B}_{n,q}^{(\alpha)}(x, y) &= \sum_{j=0}^n \binom{mx}{j} j! \sum_{k=0}^{n-j} \binom{n}{k}_q m^{j-n} \mathfrak{B}_{k,q}^{(\alpha)}(0, y) S_2(n-k, j), \\ \mathfrak{E}_{n,q}^{(\alpha)}(x, y) &= \sum_{j=0}^n \binom{mx}{j} j! \sum_{k=0}^{n-j} \binom{n}{k}_q m^{j-n} \mathfrak{E}_{k,q}^{(\alpha)}(0, y) S_2(n-k, j). \end{aligned}$$

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equations

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equ...

The familiar q -Stirling numbers $S(n, k)$ of the second kind are defined by

$$(e_q(t) - 1)^k = \sum_{m=0}^{\infty} S_{2,q}(m, k) \frac{t^m}{[m]_q!},$$

where $k \in \mathbb{N}$. Next we give the relationship between q -Bernstein basis defined by Phillips [27] and q -Bernoulli polynomials

$$b_{n,k}(q; x) := \begin{bmatrix} n \\ k \end{bmatrix}_q x^k (1-x)_{q}^{n-k}.$$

Theorem 19 *We have*

$$b_{n,k}(q; x) = x^k \sum_{m=0}^n \begin{bmatrix} n \\ m \end{bmatrix}_q S_{2,q}(m, k) \mathfrak{B}_{n-m,q}^{(k)}(1, -x). \quad (19)$$

Proof The proof follows from the following identities:

$$\begin{aligned} \frac{x^k t^k}{[k]_q!} e_q(t) E_q(-xt) &= \frac{x^k t^k}{[k]_q!} \sum_{n=0}^{\infty} \frac{(1-x)_q^n t^n}{[n]_q!} = \sum_{n=k}^{\infty} \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{x^k (1-x)_q^{n-k} t^n}{[n]_q!} \\ &= \sum_{n=k}^{\infty} b_{n,k}(q; x) \frac{t^n}{[n]_q!} \end{aligned}$$

and

$$\begin{aligned} \frac{x^k t^k}{[k]_q!} e_q(t) E_q(-xt) &= \frac{x^k (e_q(t) - 1)^k}{[k]_q!} \frac{t^k}{(e_q(t) - 1)^k} e_q(t) E_q(-xt) \\ &= x^k \sum_{m=0}^{\infty} S_{2,q}(m, k) \frac{t^m}{[m]_q!} \sum_{n=0}^{\infty} \mathfrak{B}_{n,q}^{(k)}(1, -x) \frac{t^n}{[n]_q!} \\ &= x^k \sum_{n=0}^{\infty} \left(\sum_{m=0}^n \begin{bmatrix} n \\ m \end{bmatrix}_q S_{2,q}(m, k) \mathfrak{B}_{n-m,q}^{(k)}(1, -x) \right) \frac{t^n}{[n]_q!}. \quad \square \end{aligned}$$

Finally, in their limit case when $q \rightarrow 1^-$, this last result (19) would reduce to the following formula for the classical Bernoulli polynomials $B_n^{(k)}(x)$ and the Bernstein basis $b_{n,k}(x) = \binom{n}{k} x^k (1-x)^{n-k}$:

$$b_{n,k}(x) = x^k \sum_{m=0}^n \binom{n}{m} S_2(m, k) \mathfrak{B}_{n-m}^{(k)}(1-x).$$

Competing interests

The author declares that he has no competing interests.

Acknowledgements

Dedicated to Professor Hari M. Srivastava. The author sincerely thanks the anonymous referees for their valuable suggestions and comments which have helped improve this article.

Received: 31 January 2013

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · *Advances in Difference Equations*

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equ...

References

1. Andrews, GE, Askey, R, Roy, R: Special Functions. Encyclopedia of Mathematics and Its Applications, vol. 71. Cambridge University Press, Cambridge (1999)
2. Carlitz, L: q -Bernoulli numbers and polynomials. *Duke Math. J.* **15**, 987-1000 (1948)
3. Carlitz, L: q -Bernoulli and Eulerian numbers. *Trans. Am. Math. Soc.* **76**, 332-350 (1954)
4. Carlitz, L: Expansions of q -Bernoulli numbers. *Duke Math. J.* **25**, 355-364 (1958)
5. Cenkci, M, Can, M: Some results on q -analogue of the Lerch zeta function. *Adv. Stud. Contemp. Math.* **12**, 213-223 (2006)
6. Cenkci, M, Can, M, Kurt, V: q -Extensions of Genocchi numbers. *J. Korean Math. Soc.* **43**, 183-198 (2006)
7. Cenkci, M, Kurt, V, Rim, SH, Simsek, Y: On $(i; q)$ -Bernoulli and Euler numbers. *Appl. Math. Lett.* **21**, 706-711 (2008)
8. Choi, J, Anderson, PJ, Srivastava, HM: Some q -extensions of the Apostol-Bernoulli and the Apostol-Euler polynomials of order n , and the multiple Hurwitz zeta function. *Appl. Math. Comput.* **199**, 723-737 (2008)
9. Choi, J, Anderson, PJ, Srivastava, HM: Carlitz's q -Bernoulli and q -Euler numbers and polynomials and a class of q -Hurwitz zeta functions. *Appl. Math. Comput.* **215**, 1185-1208 (2009)
10. Kim, T: Some formulae for the q -Bernoulli and Euler polynomial of higher order. *J. Math. Anal. Appl.* **273**, 236-242 (2002)
11. Kim, T: q -Generalized Euler numbers and polynomials. *Russ. J. Math. Phys.* **13**, 293-298 (2006)
12. Kim, T, Kim, YH, Hwang, KW: On the q -extensions of the Bernoulli and Euler numbers, related identities and Lerch zeta function. *Proc. Jangjeon Math. Soc.* **12**, 77-92 (2009)
13. Kim, T, Rim, SH, Simsek, Y, Kim, D: On the analogs of Bernoulli and Euler numbers, related identities and zeta and L -functions. *J. Korean Math. Soc.* **45**, 435-453 (2008)
14. Ozden, H, Simsek, Y: A new extension of q -Euler numbers and polynomials related to their interpolation functions. *Appl. Math. Lett.* **21**, 934-939 (2008)
15. Ryoo, CS, Seo, JJ, Kim, T: A note on generalized twisted q -Euler numbers and polynomials. *J. Comput. Anal. Appl.* **10**, 483-493 (2008)
16. Simsek, Y: q -Analogue of the twisted L -series and q -twisted Euler numbers. *J. Number Theory* **110**, 267-278 (2005)
17. Simsek, Y: Twisted $(h; q)$ -Bernoulli numbers and polynomials related to twisted $(h; q)$ -zeta function and L -function. *J. Math. Anal. Appl.* **324**, 790-804 (2006)
18. Simsek, Y: Generating functions of the twisted Bernoulli numbers and polynomials associated with their interpolation functions. *Adv. Stud. Contemp. Math.* **16**, 251-278 (2008)
19. Luo, QM, Srivastava, HM: q -Extensions of some relationships between the Bernoulli and Euler polynomials. *Taiwan. J. Math.* **15**, 241-257 (2011)
20. Srivastava, HM, Kim, T, Simsek, Y: q -Bernoulli numbers and polynomials associated with multiple q -zeta functions and basic L -series. *Russ. J. Math. Phys.* **12**, 241-268 (2005)
21. Srivastava, HM: Some generalizations and basic (or q -) extensions of the Bernoulli, Euler and Genocchi polynomials. *Appl. Math. Inf. Sci.* **5**, 390-444 (2011)
22. Mahmudov, NI: q -Analogues of the Bernoulli and Genocchi polynomials and the Srivastava-Pintér addition theorems. *Discrete Dyn. Nat. Soc.* **2012**, Article ID 169348 (2012). doi:10.1155/2012/169348
23. Chan, OY, Manna, D: A new q -analogue for Bernoulli numbers. Preprint. oyeat.com/papers/qBernoulli-20110825.pdf
24. Srivastava, HM, Pintér, Á: Remarks on some relationships between the Bernoulli and Euler polynomials. *Appl. Math. Lett.* **17**, 375-380 (2004)
25. Cheon, GS: A note on the Bernoulli and Euler polynomials. *Appl. Math. Lett.* **16**, 365-368 (2003)
26. Luo, QM: Some results for the q -Bernoulli and q -Euler polynomials. *J. Math. Anal. Appl.* **363**, 7-18 (2010)
27. Phillips, GM: On generalized Bernstein polynomials. In: *Numerical Analysis*, pp. 263-269. World Scientific, River Edge (1996)

doi:10.1186/1687-1847-2013-108

Cite this article as: Mahmudov: On a class of q -Bernoulli and q -Euler polynomials. *Advances in Difference Equations* 2013 **2013**:108.

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equations

Citations 13

References 34

Determinant approach to the 2-Iterated q-Appell and mixed type q-Appell polynomials

" $G_n, q := G_n, q(0)$ [7, 8] Al-Salaam showed that the class of all q-Appell polynomials is a maximal commutative subgroup of the group of all polynomial sets, i.e. the class of all q-Appell sequences is closed under the operation of q-umbral composition of polynomial sequences. "

[\[Show abstract\]](#)

No preview · Article · Jun 2016

 Subuhi Khan  Mumtaz Riyasat

[Read](#)

0 Comment 0 Citation

A Note on the Apostol Type q-Frobenius-Euler Polynomials and Generalizations of the Srivastava-Pinter Addition Theorems

"Choi et al.[6] investigated q-Euler and q-Bernoulli polynomials and gave the relation between these polynomials, Choi et al.[7] proved some relation for the Apostol-Euler polynomials and Apostol-Bernoulli polynomials, Srivastava et al. [27] proved some relation for q-Bernoulli polynomials and multiple q-Zeta function. Finally, Mahmudov in ([16], [17]) by using q-quantum calculus, introduced and gave some relations for the q-Bernoulli polynomials and q-Euler polynomials with two variable x, y. In this work, we introduce q-Apostol type Frobenius-Euler polynomials. "

[\[Show abstract\]](#)

No preview · Article · Jan 2016 · Filomat

 Burak Kurt

[Read](#)

0 Comment 0 Citation

Una nueva clase de polinomios q-Apostol-Bernoulli de orden α

"Bernoulli numbers were first studied by Carlitz[3]. Thereafter various other analogues of Bernoulli numbers and polynomials have been studied arising from varying motivations. Many authors have further studied and developed this subject, among which a few to mention are Koblitz [11], Tsumura [26], Srivastava et al. [24], Cenkciand Can[4], Ernst [7], Ryo [22], Choi et al. [5], Kim et al. [10], Luo [17], Luo and Srivastava [15], Mahmudov [19] and Lee and Ryo[13]. We recall here some of these definitions. "

[\[Show abstract\]](#)

Full-text · Article · Jul 2014

 Mridula Garg  Subhash Alha

[Read full-text](#)

People who read this publication also read:

Article: On a class of generalized q-Bernoulli and q-Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equ...

Note: This list is based on the publications in our de

Similar publications

On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Article · Apr 2013 · Advances in Difference Equations

 Nazim I Mahmudov  M. Eini Keleshteri

[Read full-text](#)

Identities involving values of Bernstein, q -Bernoulli and q -Euler polynomials

Full-text · Article · Jan 2011 · Russian Journal of Mathematical Physics

 A. Bayad  T Kim

[Read full-text](#)

Some identities of higher order Barnes-type q -Bernoulli polynomials and higher order Barnes-type q -Euler polynomials

Full-text · Article · Dec 2015 · Advances in Difference Equations

 Lee-Chae Jang  Sang-Ki Choi  Hyuck In Kwon

[Read full-text](#)

Data provided are for informational purposes only. Although carefully collected, accuracy cannot be guaranteed. The impact factor represents a rough estimation of the journal's impact factor and does not reflect the actual current impact factor. Publisher conditions are provided by RoMEO. Differing provisions from the publisher's actual policy or licence agreement may be applicable.

[Show self-archiving restrictions](#)

Last Updated: 13 Jan 16

© 2008-2016 researchgate.net. All rights reserved.

[About us](#) · [Contact us](#) · [Careers](#) · [Developers](#) · [News](#) · [Help Center](#) · [Privacy](#) · [Terms](#) | [Advertising](#) · [Recruiting](#)

People who read this publication also read:

Article: On a class of generalized q -Bernoulli and q -Euler polynomials

Full-text · Apr 2013 · Advances in Difference Equ...
