

Bernoulli numbers and polynomials from a more general point of view

G. DATTOLI – C. CESARANO – S. LORENZUTTA

RIASSUNTO: *Si applica il metodo della funzione generatrice per introdurre nuove forme di numeri e polinomi di Bernoulli che vengono utilizzati per sviluppare e calcolare somme parziali che coinvolgono polinomi a più indici e a più variabili. Si sviluppano considerazioni analoghe per i polinomi ed i numeri di Eulero.*

ABSTRACT: *We apply the method of generating function, to introduce new forms of Bernoulli numbers and polynomials, which are exploited to derive further classes of partial sums involving generalized many index many variable polynomials. Analogous considerations are developed for the Euler numbers and polynomials.*

1 – Introduction

In a previous paper [1] we have derived partial sums involving Hermite, Laguerre and Appell polynomials in terms of generalized Bernoulli polynomials. The new and interesting possibilities offered by this class of polynomials are better illustrated by an example relevant to the derivations of a partial sum involving two-index Hermite polynomials.

To this aim we remind that:

$$(1) \quad \sum_{n=0}^{N-1} (x + ny)^r = \frac{y^r}{r+1} \left[B_{r+1} \left(N + \frac{x}{y} \right) - B_{r+1} \left(\frac{x}{y} \right) \right]$$

KEY WORDS AND PHRASES: *Monomiality – Polynomials – Hermite – Bernoulli – Euler – Partial sums.*

A.M.S. CLASSIFICATION: 33C25

where $B_n(x)$ are Bernoulli polynomials defined by the generating function:

$$(2) \quad \frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} \frac{t^n}{n!} B_n(x)$$

or in terms of Bernoulli numbers B_n as:

$$(3) \quad B_n(x) = \sum_{s=0}^n \binom{n}{s} B_{n-s} x^s .$$

Hermite polynomials with two variables and one parameter can be defined by means of the operational identity [2]:

$$(4) \quad e^{\tau \frac{\partial^2}{\partial x \partial y}} \{x^m y^n\} = h_{m,n}(x, y | \tau),$$

and the $h_{m,n}(x, y | \tau)$ are defined by the double sum:

$$(5) \quad h_{m,n}(x, y | \tau) = m!n! \sum_{s=0}^{\min(m,n)} \frac{\tau^s x^{m-s} y^{n-s}}{s!(m-s)!(n-s)} .$$

The identity (4) can be used to state that:

$$(6) \quad e^{\tau \frac{\partial^2}{\partial x \partial y}} \{(ax + b)^m (cy + d)^n\} = h_{m,n}(ax + b, cy + d | ac\tau),$$

and to introduce the following two variable one parameter Bernoulli polynomials:

$$(7) \quad {}_h B_{r,s}(x, y | \tau) = \sum_{q=0}^r \sum_{k=0}^s \binom{r}{q} \binom{s}{k} B_{r-q} B_{s-k} h_{q,k}(x, y | \tau) .$$

It is easy to note that from equation (4) we get:

$$(8) \quad e^{\tau \frac{\partial^2}{\partial x \partial y}} \{B_r(x) B_s(y)\} = {}_h B_{r,s}(x, y | \tau) .$$

This new class of Bernoulli polynomials can be used to derive the following important result:

$$(9) \quad \sum_{m=0}^{M-1} \sum_{n=0}^{N-1} h_{r,s}(x+my, z+nw | \tau) = \frac{y^r w^s}{(r+1)(s+1)} \times \\ \times \left\{ {}_h B_{r+1,s+1} \left(M + \frac{x}{y}, N + \frac{z}{w} \middle| \frac{\tau}{yw} \right) - {}_h B_{r+1,s+1} \left(M + \frac{x}{y}, \frac{z}{w} \middle| \frac{\tau}{yw} \right) + \right. \\ \left. - {}_h B_{r+1,s+1} \left(\frac{x}{y}, N + \frac{z}{w}, \frac{\tau}{yw} \right) + {}_h B_{r+1,s+1} \left(M + \frac{x}{y}, \frac{z}{w} \middle| \frac{\tau}{yw} \right) \right\}$$

which is a consequence of equations (1), (6), (7), (8). The generating function of this last class of Bernoulli polynomials can be shown to be provided by:

$$(10) \quad \frac{uv e^{ux+vy+\tau uv}}{(e^u - 1)(e^v - 1)} = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{u^m}{m!} \frac{v^n}{n!} ({}_h B_{m,n}(x, y | \tau)).$$

This introductory examples shows that a wealth of implications is offered by the use of generalized forms of Bernoulli polynomials. In the forthcoming sections we will develop a more systematic analysis which yields a deeper insight into the effectiveness of this type of generalizations.

2 – Finite sums and new classes of Bernoulli numbers

In ref. [1] we have touched on the following new class of numbers:

$$(11) \quad {}_H B_n = \sum_{s=0}^{[n/2]} \frac{n! B_{n-2s} B_s}{s!(n-2s)!}$$

which are recognized as an Hermite convolution of Bernoulli numbers on themselves. The generating functions of the ${}_H B_n$ is provided by:

$$(12) \quad \frac{t^3}{(e^t - 1)(e^{t^2} - 1)} = \sum_{n=0}^{\infty} \frac{t^n}{n!} ({}_H B_n)$$

which suggests the following generalization ($(a, b) \neq 0$):

$$(13) \quad \frac{t^3}{(e^{at} - 1)(e^{bt^2} - 1)} = \sum_{n=0}^{\infty} \frac{t^n}{n!} ({}_H B_n^*(a, b))$$

with:

$$(14) \quad {}_H B_n^*(a, b) = \frac{1}{ab} \sum_{s=0}^{[n/2]} \frac{n! a^{n-2s} b^s B_{n-2s} B_s}{s!(n-2s)!}.$$

It is also evident that the generating function:

$$(15) \quad \frac{t^3 e^{xt+yt^2}}{(e^{at}-1)(e^{bt^2}-1)} = \sum_{n=0}^{\infty} \frac{t^n}{n!} ({}_H B_n^*(a, b | x, y))$$

can be exploited to define the polynomials:

$$(16) \quad {}_H B_n^*(a, b | x, y) = \sum_{s=0}^n \binom{n}{s} ({}_H B_{n-s}^*(a, b) H_s(x, y))$$

as results from equations (13) and (15) which are also expressed as:

$$(17) \quad {}_H B_n^*(a, b | x, y) = \frac{n!}{ab} \sum_{s=0}^{[n/2]} \frac{n! a^{n-2s} b^s}{s!(n-2s)!} B_{n-2s} \left(\frac{x}{a}\right) B_s \left(\frac{y}{b}\right)$$

yielding:

$$(18) \quad {}_H B_n^*(a, b | 0, 0) = {}_H B_n^*(a, b)$$

where $H_n(x, y) = n! \sum_{s=0}^{[n/2]} \frac{y^s x^{n-2s}}{s!(n-2s)!}$ are the Kampè de Fèriet polynomials. The use of polynomials (16) is suggested by partial sums of the type:

$$(19) \quad \sum_{m=0}^{M-1} \sum_{n=0}^{N-1} H_r(x+my, z+nw) = \frac{1}{(r+1)(r+2)(r+3)} \times \\ \times \left\{ {}_H B_{r+3}^*(y, w | x+My, z+Nw) - {}_H B_{r+3}^*(y, w | x+My, z) + \right. \\ \left. - {}_H B_{r+3}^*(y, w | x, z+Nw) + {}_H B_{r+3}^*(y, w | x, z) \right\}.$$

which provides one of the main results of the paper.

A further application of polynomials ${}_H B_n^*(a, b \mid x, y)$ is relevant to the multiplication theorems. We find indeed:

$$(20) \quad \begin{aligned} & {}_H B_n(m x, m^2 y) = \\ & = \frac{m^{n-2}}{(n+1)(n+2)} \sum_{k=0}^{m-1} \sum_{h=0}^{m^2-1} \left({}_H B_{n+2}^* \left(1, 1 \mid x + \frac{k}{m}, y + \frac{h}{m^2} + \frac{1}{m^2} \right) \right) + \\ & \quad - \left({}_H B_{n+2}^* \left(1, 1 \mid x + \frac{k}{m}, y + \frac{h}{m^2} \right) \right) \end{aligned}$$

and

$$(21) \quad {}_H B_n^*(a, b \mid m x, m^2 y) = m^{n-3} \sum_{k=1}^{m-1} \sum_{h=1}^{m^2-1} \left({}_H B_n^* \left(a, b \mid x + \frac{a k}{m}, y + \frac{b h}{m^2} \right) \right)$$

and:

$$(22) \quad {}_h B_{m,n}(p x, q y \mid \tau) = p^{m-1} q^{n-1} \sum_{k=0}^{p-1} \sum_{h=0}^{q-1} \left({}_h B_{m,n} \left(x + \frac{k}{p}, y + \frac{h}{q} \mid \frac{\tau}{pq} \right) \right)$$

can be proved by exploiting the procedure outlined in appendix.

3 – Euler polynomials

The examples we have provided yields an idea of the implications offered by this type of generalization.

It is also evident that the considerations we have developed for Bernoulli polynomials can be extended to Euler polynomials [3]:

$$(23) \quad \frac{2e^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} \frac{t^n}{n!} E_n(x).$$

In analogy of ref.[1] and the results of previous sections, we introduce the following classes of Euler polynomials:

$$(24) \quad \frac{2e^{xt+yt^2}}{e^t + 1} = \sum_{n=0}^{\infty} \frac{t^n}{n!} ({}_H E_n(x, y))$$

and:

$$(25) \quad \frac{2^2 e^{xu+yv+\tau uv}}{(e^u+1)(e^v+1)} = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{u^m}{m!} \frac{v^n}{n!} ({}_h E_{m,n}(x, y | \tau))$$

It is easily realized that:

$$(26) \quad {}_H E_n(x, y) = \sum_{s=0}^{[n/2]} \frac{y^s E_{n-2s}(x)}{s!(n-2s)!}$$

and:

$$(27) \quad {}_h E_{m,n}(x, y | \tau) = \sum_{s=0}^m \sum_{r=0}^n \binom{m}{s} \binom{n}{r} E_{m-s} E_{n-r} h_{s,r}(x, y | \tau)$$

where $E_n(x)$ are the ordinary Euler polynomials and that the following theorems hold:

$$(28) \quad {}_H E_n(x+z, y+w) = \sum_{s=0}^n \binom{n}{s} ({}_H E_{n-s}(x, y)) H_s(z, w)$$

and (see appendix):

$$(29) \quad {}_H E_n(mx, py) = m^n \sum_{k=0}^{m-1} (-1)^k ({}_H E_n(x + \frac{k}{m}, \frac{py}{m^2})) .$$

As to the polynomials ${}_h E_{m,n}(x, y | \tau)$ we can also state that:

$$(30) \quad {}_h E_{m,n}(x+z, y+w | \tau) = \sum_{s=0}^m \sum_{r=0}^n \binom{m}{s} \binom{n}{r} ({}_h E_{m-s, n-r}(x, y | \tau)) z^s w^r$$

and:

$$(31) \quad {}_h E_{m,n}(px, qy | \tau) = p^m q^n \sum_{k=0}^{p-1} \sum_{h=0}^{q-1} (-1)^{k+h} {}_h E_{m,n}(x + \frac{k}{p}, y + \frac{h}{q} | \frac{\tau}{pq}) .$$

4 – Concluding remarks

The introduction of the Hermite-Euler polynomials given by equation (24) offers the possibility of speculating about alternative definitions as e.g. ($t < \sqrt{\pi}$):

$$(32) \quad \frac{4e^{xt+yt^2}}{(e^t + 1)(e^{t^2} + 1)} = \sum_{n=0}^{\infty} \frac{t^n}{n!} E_n^{(2)}(x, y).$$

The polynomials $E_n^{(2)}(x, y)$ are defined as:

$$(33) \quad E_n^{(2)}(x, y) = n! \sum_{s=0}^{[n/2]} \frac{E_s(y) E_{n-2s}(x)}{s!(n-2s)!}$$

and are shown to satisfy the following differential equation:

$$(34) \quad \frac{\partial}{\partial y} E_n^{(2)}(x, y) = \frac{\partial^2}{\partial x^2} E_n^{(2)}(x, y)$$

and the multiplication formula:

$$(35) \quad E_n^{(2)}(mx, m^2y) = m^n \sum_{k=0}^{m-1} \sum_{h=0}^{m^2-1} (-1)^{k+h} \left(E_n^{(2)} \left(x + \frac{k}{m}, y + \frac{h}{m^2} \right) \right).$$

The more general theorem relevant to $E_n^{(2)}(mx, py)$ requires the introduction of a class of Euler polynomials analogous to those provided by equation (13) for the Bernoulli case.

Before concluding this paper we want to emphasize that most of the identities holding for the ordinary Bernoulli or Euler polynomials can be extended to the generalized case. For example the identity [4]:

$$(36) \quad B_n(x+1) - B_n(x) = nx^{n-1}$$

can be generalized as:

$$(37) \quad {}_H B_n(x+1, y) - {}_H B_n(x, y) = nH_{n-1}(x, y)$$

which is a consequence of:

$$(38) \quad e^{y \frac{\partial^2}{\partial x^2}} B_n(x) = {}_H B_n(x, y).$$

For analogous reasons, we find:

$$(39) \quad {}_h B_{m,n}(x+1, y+1 | \tau) - {}_h B_{m,n}(x, y+1 | \tau) - {}_h B_{m,n}(x+1, y | \tau) + {}_h B_{m,n}(x, y | \tau) = mn h_{m-1, n-1}(x, y | \tau)$$

and:

$$(40) \quad {}_H B_n^*(a, b | x+a, y+b) - {}_H B_n^*(a, b | x, y) = n(n-1)(n-2) H_{n-3}(x, y)$$

and

$$(41) \quad {}_H E_n^{(2)}(x+1, y) + {}_H E_n^{(2)}(x, y) = 2H_n(x, y).$$

We can also define the further generalized form:

$$(42) \quad \prod_{i=1}^N \frac{t^{i+1} e^{x_i t^i}}{(e^{a_i t^i} - 1)} = \sum_{n=0}^{+\infty} \frac{t^n}{n!} ({}_H B_n^*(\{a_i\} | \{x_i\}))$$

thus getting ($n > N$):

$$(43) \quad {}_H B_n^*(\{a_i\} | \{x_i\}) - {}_H B_n^*(\{a_i\} | \{x_i\}) = \frac{n!}{(n-N)!} H_{n-N}(\{x_i\})$$

with:

$$(44) \quad \sum_{n=0}^{+\infty} \frac{t^n}{n!} H_n(\{x_i\}) = e^{\sum_{i=1}^N x_i t^i}.$$

The results of this paper show that the combination of operational rules and the properties of ordinary and generalized polynomials offer a wealth of possibilities to introduce new families of Euler and Bernoulli polynomials which provides a powerful tool in applications.

– Appendix

The multiplication formulae are easily stated by exploiting the method of the generating functions and suitable manipulations. We note indeed that:

$$(A.1) \quad \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{u^m}{m!} \frac{v^n}{n!} ({}_h B_{m,n}(px, qy | \tau)) = \frac{uve^{upx+vyq+\tau uv}}{(e^u - 1)(e^v - 1)}.$$

The r.h.s. of the above relation can be more conveniently rewritten as:

$$(A.2) \quad \frac{uve^{upx+vyq+\tau uv}}{(e^u - 1)(e^v - 1)} = \frac{(up)(vq)}{pq} \frac{uve^{upx+vyq+uvpq\frac{\tau}{pq}}}{(e^{up} - 1)(e^{vq} - 1)} \frac{(e^{up} - 1)(e^{vq} - 1)}{(e^u - 1)(e^v - 1)}$$

by noting that:

$$(A.3) \quad \frac{e^{up} - 1}{e^u - 1} = \sum_{r=0}^{p-1} e^{ru}$$

we can rearrange (A.2) as:

$$(A.4) \quad \frac{uve^{upx+vyq+\tau uv}}{(e^u - 1)(e^v - 1)} = \sum_{k=0}^{p-1} \sum_{h=0}^{q-1} \frac{1}{pq} \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(up)^m}{m!} \frac{(vq)^n}{n!} \times \\ \times {}_h B_{m,n}\left(x + \frac{k}{p}, y + \frac{h}{q} \middle| \frac{\tau}{pq}\right)$$

which once confronted with (A.1) yields equation (22). A similar procedure can be exploited to prove the multiplication formulae relevant to the Euler's generalized forms.

We note indeed:

$$(A.5) \quad \sum_{n=0}^{\infty} \frac{t^n}{n!} ({}_H E_n(mx, py)) = \frac{2e^{mxt+pyt^2}}{e^t + 1}$$

and handling the r.h.s. of the above equations, we find:

$$(A.6) \quad \sum_{n=0}^{\infty} \frac{t^n}{n!} ({}_H E_n(mx, py)) = \frac{2e^{mxt}}{e^{mt} + 1} \frac{e^{mt} + 1}{e^t + 1} e^{pyt^2}.$$

By noting that:

$$(A.7) \quad \frac{2e^{mxt}}{e^{mt} + 1} \frac{e^{mt} + 1}{e^t + 1} e^{pyt^2} = \sum_{k=0}^{m-1} (-1)^k \sum_{q=0}^{\infty} \frac{t^q m^q}{q!} E_q \left(x + \frac{k}{m} \right) \sum_{r=0}^{\infty} \frac{t^{2r} p^r}{r!} y^r$$

we obtain:

$$(A.8) \quad \sum_{n=0}^{\infty} \frac{t^n}{n!} ({}_H E_n(mx, py)) = \sum_{n=0}^{\infty} t^n m^n \sum_{k=0}^{m-1} (-1)^k \sum_{r=0}^{\lfloor n/2 \rfloor} \frac{E_{n-2r} \left(x + \frac{k}{m} \right)}{(n-2r)! r!} \left(\frac{py}{m^2} \right)^r$$

and using the (26) we finally state the relation (29).

REFERENCES

- [1] G. DATTOLI – S. LORENZUTTA – C. CESARANO: *Finite sums and generalization forms of Bernoulli polynomials*, Rend. di Matematica, **19** (1999), 385-391.
- [2] G. DATTOLI – A. TORRE: *Theory and applications of generalized Bessel functions*, ARACNE Editrice, Roma, 1996.
- [3] L.M. MILNE-THOMSON: *Calculus of finite differences*, MacMillan & C. Ltd., London, 1951.
- [4] M. ABRAMOVITZ – I. STEGUN: *Hand book of Mathematical Functions*, Dover Pub. Inc. New York, 1970.

*Lavoro pervenuto alla redazione il 7 marzo 2001
ed accettato per la pubblicazione il 16 luglio 2002.
Bozze licenziate il 15 gennaio 2003*

INDIRIZZO DEGLI AUTORI:

G. Dattoli – ENEA – Dipartimento Innovazione – Divisione Fisica Applicata – Centro Ricerche Frascati – C.P. 65 - 00044 Frascati – Roma (Italia)

C. Cesarano – University of ULM – Department of Mathematics - ULM - Germany and ENEA Guest

E-mail: cesarano@frascati.enea.it

S. Lorenzutta – ENEA – Dipartimento Innovazione – Divisione Fisica Applicata – Centro Ricerche Bologna – Via Don Fiammelli – Bologna (Italia)