

# A survey on the theory of multiple Bernoulli polynomials and multiple $L$ -functions of root systems

By

Yasushi KOMORI\*, Kohji MATSUMOTO\*\* and Hirofumi TSUMURA\*\*\*

## § 1. Introduction

In [36], Witten found that a certain series of Dirichlet type appear in two dimensional quantum gauge theories with connected compact semisimple Lie groups. Motivated by this observation, Zagier [37] defined the Witten zeta-functions as

$$(1.1) \quad \zeta_W(s; \mathfrak{g}) = \sum_{\varphi} \frac{1}{(\dim \varphi)^s}$$

for  $s \in \mathbb{C}$ , where the summation runs over all finite dimensional irreducible representations  $\varphi$  of a given semisimple Lie algebra  $\mathfrak{g}$ . It is known that a semisimple Lie algebra is a direct sum of simple Lie algebras and simple Lie algebras of rank  $r$  are associated to an irreducible root system of type  $X_r$  where  $X = A, B, \dots, G$  (see Section 4 for the details). In the case where  $\mathfrak{g}$  is of type  $A_1$ , the series reduces to the Riemann zeta-function

$$(1.2) \quad \zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}.$$

It is well known that the notion of “zeta-functions” plays an important tool in various areas of modern mathematics.

When  $s$  is an even positive integer, then their values are crucial (if  $s$  is an odd integer, those with appropriate characters play the same role); mathematically, they

---

Received April 20, 201x. Revised September 11, 201x.

2000 Mathematics Subject Classification(s): 11M41, 22E70

\*Department of Mathematics, Rikkyo University, Nishi-Ikebukuro, Toshima-ku, Tokyo 171-8501, Japan.

e-mail: komori@rikkyo.ac.jp

\*\*Graduate School of Mathematics, Nagoya University, Chikusa-ku, Nagoya 464-8602, Japan.

e-mail: kohjimat@math.nagoya-u.ac.jp

\*\*\*Department of Mathematics and Information Sciences, Tokyo Metropolitan University, 1-1, Minami-Ohsawa, Hachioji, Tokyo 192-0397, Japan.

e-mail: tsumura@tmu.ac.jp

give the volumes of certain moduli spaces of flat connections, and physically, the 0-th orders of the partition functions of two dimensional quantum gauge theories. Assume that  $s$  is an even positive integer  $2k$ . Witten and Zagier showed that their values are in  $\mathbb{Q}\pi^{|\Delta_+|2k}$ , where  $\Delta_+$  denotes the set of all positive roots. Euler already evaluated them in the  $A_1$  case. The  $A_2$  case was first studied by Tornheim [33] and Mordell [29] independently, and further considered by several authors [7, 31, 34]. In [32], Szenes gave a certain algorithm for the computation in general cases, from the viewpoint of hyperplane arrangements. Gunnells and Sczech also gave another general algorithm and the explicit forms in the  $A_3$  case as an application [6].

This article is a survey on a new approach to this problem proposed in [11, 12, 15–18, 22, 28] and is an extended and updated version of the informal articles [13, 14]. We will introduce generalizations of Bernoulli polynomials and zeta-functions associated with root systems, which include the Riemann zeta-function, the Euler-Zagier zeta-functions and the Witten zeta-functions. Furthermore we will develop a theory similar to that of the classical Riemann zeta-function.

Acknowledgment: The authors would like to thank the referee for critical reading of the manuscript and useful comments.

## § 2. A Review of Classical Theory

Before stating our results, first we recall the classical theory of the Riemann zeta-function and Bernoulli numbers.

The following is a well-known formula for the Riemann zeta-function and Bernoulli numbers: For  $k \in \mathbb{Z}_{\geq 1}$ ,

$$(2.1) \quad 2\zeta(2k) = -B_{2k} \frac{(2\pi i)^{2k}}{(2k)!},$$

where the definition of  $B_k$  is given by, for  $t \in \mathbb{C}$  with  $|t| < 2\pi$ ,

$$(2.2) \quad \frac{t}{e^t - 1} = \sum_{k=0}^{\infty} B_k \frac{t^k}{k!}.$$

Using this formula, we obtain for  $k \in \mathbb{Z}_{\geq 1}$ ,

$$(2.3) \quad \zeta(2k) + (-1)^{2k} \zeta(2k) = -B_{2k} \frac{(2\pi i)^{2k}}{(2k)!},$$

$$(2.4) \quad \zeta(2k+1) + (-1)^{2k+1} \zeta(2k+1) = -B_{2k+1} \frac{(2\pi i)^{2k+1}}{(2k+1)!} = 0.$$

Hence we have the following important relations: For  $k \in \mathbb{Z}_{\geq 2}$ ,

$$(2.5) \quad \zeta(k) + (-1)^k \zeta(k) = -B_k \frac{(2\pi i)^k}{k!},$$

that is, value-relations can be written in terms of Bernoulli numbers.

These relations are generalized to the case of Lerch zeta-functions and periodic Bernoulli functions. Let  $\varphi(s, y)$  be the Lerch zeta-function defined by

$$(2.6) \quad \varphi(s, y) = \sum_{n=1}^{\infty} \frac{e^{2\pi i n y}}{n^s}.$$

Then a formula for Lerch zeta-functions implies that for  $k \in \mathbb{Z}_{\geq 2}$  and  $y \in \mathbb{R}$ ,

$$(2.7) \quad \varphi(k, y) + (-1)^k \varphi(k, -y) = -B_k(\{y\}) \frac{(2\pi i)^k}{k!},$$

that is, functional relations as functions in  $y$  can be written in terms of periodic Bernoulli functions, which are defined by

$$(2.8) \quad \frac{te^{t\{y\}}}{e^t - 1} = \sum_{k=0}^{\infty} B_k(\{y\}) \frac{t^k}{k!},$$

and  $\{y\} = y - [y]$  is the fractional part of  $y$ .

Once we obtain the notion of periodic Bernoulli functions, we can calculate special values of Dirichlet  $L$ -functions  $L(s, \chi)$  in terms of them. For a primitive character  $\chi$  of conductor  $f$  and  $k \in \mathbb{Z}_{\geq 2}$  satisfying  $(-1)^k \chi(-1) = 1$ , we have

$$(2.9) \quad L(k, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^k} = \frac{(-1)^{k+1}}{2} \frac{(2\pi i)^k}{k! f^k} g(\chi) B_{k, \bar{\chi}},$$

where  $g(\chi)$  is the Gauss sum,  $\bar{\chi}$  is the complex conjugate of  $\chi$ , and

$$(2.10) \quad B_{k, \chi} = f^{k-1} \sum_{a=1}^f \chi(a) B_k(a/f).$$

Our aim is to find a good class of multiple zeta-functions which generalizes the theory above. First we will introduce zeta- and  $L$ -functions associated with semisimple Lie algebras, which are corresponding to simply-connected Lie groups. Moreover besides those, we will study zeta-functions associated with Lie groups that may not be simply-connected.

### § 3. An Overview of Our Results

Based on the observation given in the previous section, we will construct multiple generalizations of Bernoulli polynomials and multiple zeta- and  $L$ -functions associated

with arbitrary root systems. Before introducing the general theory, we give two simple theorems without using the terminology of root systems. For  $s_1, s_2, s_3 \in \mathbb{C}$  with  $\Re s_1, \Re s_2, \Re s_3 \geq 2$  and  $y_1, y_2 \in \mathbb{R}$ , we consider the convergent series

$$(3.1) \quad \zeta_2(s_1, s_2, s_3, y_1, y_2; A_2) = \sum_{m, n=1}^{\infty} \frac{e^{2\pi i(m y_1 + n y_2)}}{m^{s_1} n^{s_2} (m+n)^{s_3}}.$$

**Theorem A.** For  $k_1, k_2, k_3 \in \mathbb{Z}_{\geq 2}$ ,

$$(3.2) \quad \begin{aligned} & \zeta_2(k_1, k_2, k_3, y_1, y_2; A_2) + (-1)^{k_1} \zeta_2(k_1, k_3, k_2, -y_1 + y_2, y_2; A_2) \\ & + (-1)^{k_2} \zeta_2(k_3, k_2, k_1, y_1, y_1 - y_2; A_2) + (-1)^{k_2+k_3} \zeta_2(k_3, k_1, k_2, -y_1 + y_2, -y_1; A_2) \\ & + (-1)^{k_1+k_3} \zeta_2(k_2, k_3, k_1, -y_2, y_1 - y_2; A_2) + (-1)^{k_1+k_2+k_3} \zeta_2(k_2, k_1, k_3, -y_2, -y_1; A_2) \\ & = (-1)^3 \mathcal{P}(k_1, k_2, k_3, y_1, y_2; A_2) \frac{(2\pi i)^{k_1+k_2+k_3}}{k_1! k_2! k_3!}, \end{aligned}$$

where  $\mathcal{P}(k_1, k_2, k_3, y_1, y_2; A_2)$  is a multiple periodic Bernoulli function (defined later). In particular, we have

$$(3.3) \quad \zeta_2(2, 2, 2, 0, 0; A_2) = \frac{1}{6} (-1)^3 \frac{1}{3780} \frac{(2\pi i)^{2+2+2}}{2!2!2!} = \frac{\pi^6}{2835}.$$

This should be compared with (2.7) and

$$(3.4) \quad \zeta(2) = \frac{1}{2} (-1) \frac{1}{6} \frac{(2\pi i)^2}{2!} = \frac{\pi^2}{6}.$$

For  $s_1, s_2, s_3 \in \mathbb{C}$  with  $\Re s_1, \Re s_2, \Re s_3 \geq 2$  and primitive Dirichlet characters  $\chi_1, \chi_2, \chi_3$ , consider the convergent series

$$(3.5) \quad L_2(s_1, s_2, s_3, \chi_1, \chi_2, \chi_3; A_2) = \sum_{m, n=1}^{\infty} \frac{\chi_1(m) \chi_2(n) \chi_3(m+n)}{m^{s_1} n^{s_2} (m+n)^{s_3}}.$$

**Theorem B.** For  $k \in \mathbb{Z}_{\geq 2}$  and a primitive Dirichlet character  $\chi$  of conductor  $f$  such that  $(-1)^k \chi(-1) = 1$ ,

$$(3.6) \quad L_2(k, k, k, \chi, \chi, \chi; A_2) = \frac{(-1)^{3k+3}}{6} \left( \frac{(2\pi i)^k}{k! f^k} g(\chi) \right)^3 B_{k, k, k, \bar{\chi}, \bar{\chi}, \bar{\chi}}(A_2),$$

where  $B_{k_1, k_2, k_3, \chi_1, \chi_2, \chi_3}(A_2)$  is a multiple generalized Bernoulli number (defined later). In particular, for the quadratic character of conductor 5, namely  $\rho_5(1) = \rho_5(4) = 1$  and  $\rho_5(2) = \rho_5(3) = -1$ , we have

$$(3.7) \quad L_2(2, 2, 2, \rho_5, \rho_5, \rho_5; A_2) = \frac{(-1)^{6+3}}{6} \left( \frac{(2\pi i)^2}{2!5^2} \sqrt{5} \right)^3 \left( -\frac{28}{125} \right) = -\frac{112\sqrt{5}}{1171875} \pi^6.$$

This should be compared with

$$(3.8) \quad \begin{aligned} L(k, \chi) &= \frac{(-1)^{k+1} (2\pi i)^k}{2} \frac{g(\chi) B_{k, \bar{\chi}}}{k! f^k}, \\ L(2, \rho_5) &= \frac{(-1)^{2+1} (2\pi i)^2}{2} \frac{\sqrt{5}}{2! 5^2} \frac{4}{5} = \frac{4\sqrt{5}}{125} \pi^2. \end{aligned}$$

Theorems A and B are special cases of our main theorems. In the following sections, we will formulate those theorems.

*Remark.* Tornheim [33] already showed that for  $a, b, c \in \mathbb{N}$ ,  $\zeta_2(a, b, c, 0, 0; A_2)$  can be expressed as a polynomial in Riemann zeta values with  $\mathbb{Q}$ -coefficients if  $a + b + c$  is odd. On the other hand, it seems difficult to treat the case when  $a + b + c$  is even. Actually, in [7], it is stated that only the following four cases can be evaluated in terms of Riemann zeta values:  $(a, b, c) = (1, 1, N - 2)$ ,  $(j, N - j - 1, 1)$ ,  $(N/3, N/3, N/3)$  and  $(N/3, N/3 - 1, N/3 + 1)$ , where  $N \in \mathbb{Z}_{\geq 3}$  is even and  $j \in \mathbb{Z}_{\geq 1}$ . Hence, for example, it is unknown whether the case  $(2p, 2q, 2r)$  can be evaluated, except for the case  $p = q = r$ . Especially, as for the double zeta value  $\zeta_2(a, 0, b, 0, 0; A_2)$  where  $a + b$  is even, it is unknown whether it can be evaluated in terms of Riemann zeta values if  $a + b \geq 8$ , except for the case  $a = b$  or  $a = 1$  (see [1, Section 4]).

## § 4. Root Systems

Now we start to describe our general theory. First, for reader's convenience, we give the definition and several examples of root systems.

### § 4.1. Definitions

Let  $V$  be an  $r$  dimensional real vector space equipped with the inner product  $\langle \cdot, \cdot \rangle$ . A root system  $\Delta \subset V$  is a set of vectors (roots) satisfying

1.  $|\Delta| < \infty$  and  $0 \notin \Delta$ ,
2.  $\sigma_\alpha \Delta = \Delta$  for all  $\alpha \in \Delta$ ,
3.  $\langle \alpha^\vee, \beta \rangle \in \mathbb{Z}$  for all  $\alpha, \beta \in \Delta$ ,
4.  $\alpha, c\alpha \in \Delta$  for  $c \in \mathbb{R} \implies c = \pm 1$ ,


where  $\sigma_\alpha$  denotes the reflection with respect to the hyperplane  $H_\alpha$  orthogonal to  $\alpha$  and  $\alpha^\vee = 2\alpha/\langle \alpha, \alpha \rangle$  (coroot). A root system  $\Delta$  is called irreducible if it cannot be partitioned into the union of two proper subsets such that each root in one set is orthogonal to each root in the other.

Let  $W$  be the Weyl group (the group generated by all  $\sigma_\alpha$ ). Let  $\{\alpha_1, \dots, \alpha_r\}$  be the set of all fundamental roots (a basis by which any  $\alpha \in \Delta$  can be written as  $\alpha = c_1\alpha_1 + \dots + c_r\alpha_r \in \Delta$ ,  $c_i \in \mathbb{Z}$  with all  $c_i \geq 0$  or all  $c_i \leq 0$ ). Let  $\Delta_+$  be the set of all positive roots (all roots  $\alpha = c_1\alpha_1 + \dots + c_r\alpha_r \in \Delta$ ,  $c_i \in \mathbb{Z}$  with all  $c_i \geq 0$ ),  $\rho = \frac{1}{2} \sum_{\alpha \in \Delta_+} \alpha$ . Let  $Q = \bigoplus_{i=1}^r \mathbb{Z}\alpha_i$  be the root lattice, Let  $P = \bigoplus_{i=1}^r \mathbb{Z}\lambda_i$  (the weight lattice) and  $P_+ = \bigoplus_{i=1}^r \mathbb{Z}_{\geq 0}\lambda_i$ , where  $\{\lambda_1, \dots, \lambda_r\}$  is the dual basis of  $\{\alpha_1^\vee, \dots, \alpha_r^\vee\}$ .

#### § 4.2. Examples

Since we mainly treat coroots in this paper, we give examples of root systems in terms of coroots. Note that if  $\Delta$  is a root system, then  $\Delta^\vee = \{\alpha^\vee \mid \alpha \in \Delta\}$  is also a root system.

There is only one root system of rank 1, that is, of type  $A_1$  and there are four root systems of rank 2, that is, of type  $A_1 \times A_1$ ,  $A_2$ ,  $C_2$  (or  $B_2$ ) and  $G_2$  (roots in the shaded region are positive):



### § 5. Zeta-Functions of Root Systems

#### § 5.1. Witten Zeta-Functions

As prototypes of zeta-functions of root systems, we give the definition of Witten zeta-functions.

**Definition 5.1** (Witten zeta-functions [36, 37]). For a complex simple Lie algebra  $\mathfrak{g}$  with the root system  $\Delta$ ,

$$(5.1) \quad \zeta_W(s; \Delta) = \sum_{\varphi} (\dim \varphi)^{-s} = K(\Delta)^s \sum_{\lambda \in P_+} \prod_{\alpha \in \Delta_+} \frac{1}{\langle \alpha^\vee, \lambda + \rho \rangle^s},$$

where the summation runs over all finite dimensional irreducible representations  $\varphi$  on the second member of the above and  $K(\Delta) \in \mathbb{Z}_{\geq 1}$  is a constant.

Note that in the second equality in Definition 5.1, we have used Weyl's dimension formula.

We also use the notation

$$(5.2) \quad \zeta_W(s; X_r) = \zeta_W(s; \Delta)$$


if  $\Delta$  is of type  $X_r$ .

**Example 5.2.** From the second expression of Definition 5.1, we obtain the explicit forms of Witten zeta-functions as follows in the  $A_1, A_2, C_2$  cases:

$$\zeta_W(s; A_1) = \sum_{m=1}^{\infty} \frac{1}{m^s} = \zeta(s),$$

$$\zeta_W(s; A_2) = 2^s \sum_{m,n=1}^{\infty} \frac{1}{m^s n^s (m+n)^s},$$

$$\zeta_W(s; C_2) = 6^s \sum_{m,n=1}^{\infty} \frac{1}{m^s n^s (m+n)^s (m+2n)^s}.$$



Comparing these with Section 4.2, we observe that each factor of the form  $am + bn$  ( $a, b \in \mathbb{Z}_{\geq 0}$ ) in the denominators corresponds to the coroot of the form  $a\alpha_1^\vee + b\alpha_2^\vee$ .

## § 5.2. Zeta-Functions of Root Systems

**Definition 5.3** (Zeta-functions of root systems [11, 15, 17, 28]). For a root system  $\Delta$ , define

$$(5.3) \quad \zeta_r(\mathbf{s}, \mathbf{y}; \Delta) = \sum_{\lambda \in P_+} e^{2\pi i \langle \mathbf{y}, \lambda + \rho \rangle} \prod_{\alpha \in \Delta_+} \frac{1}{\langle \alpha^\vee, \lambda + \rho \rangle^{s_\alpha}},$$

where  $\mathbf{s} = (s_\alpha)_{\alpha \in \Delta_+} \in \mathbb{C}^{|\Delta_+|}$  and  $\mathbf{y} \in V$ .

As in the case of Witten zeta-functions, we may write

$$(5.4) \quad \zeta_r(\mathbf{s}, \mathbf{y}; \Delta) = \zeta_r(\mathbf{s}, \mathbf{y}; X_r)$$

if  $\Delta$  is of type  $X_r$ . It is easy to see that (5.3) with  $\mathbf{y} = \mathbf{0}$  is essentially a multi-variable version of Witten zeta-functions. Indeed we see that  $\zeta_W(s; \Delta) = K(\Delta)^s \zeta_r((s, \dots, s), \mathbf{0}; \Delta)$ .

To define an action of the Weyl group, we extend  $\mathbf{s} = (s_\alpha)_{\alpha \in \Delta_+}$  to  $(s_\alpha)_{\alpha \in \Delta}$  by  $s_\alpha = s_{-\alpha}$  and define  $(w\mathbf{s})_\alpha = s_{w^{-1}\alpha}$ . Then we have our first theorem.

**Theorem 5.4** ([17]). For  $\mathbf{s} = \mathbf{k} = (k_\alpha)_{\alpha \in \Delta_+} \in \mathbb{Z}_{\geq 2}^{|\Delta_+|}$ , we have

$$(5.5) \quad \sum_{w \in W} \left( \prod_{\alpha \in \Delta_+ \cap w\Delta_-} (-1)^{k_\alpha} \right) \zeta_r(w^{-1}\mathbf{k}, w^{-1}\mathbf{y}; \Delta) = (-1)^{|\Delta_+|} \mathcal{P}(\mathbf{k}, \mathbf{y}; \Delta) \left( \prod_{\alpha \in \Delta_+} \frac{(2\pi i)^{k_\alpha}}{k_\alpha!} \right),$$

where  $\mathcal{P}(\mathbf{k}, \mathbf{y}; \Delta)$  is a multiple periodic Bernoulli function (defined later).

**Example 5.5.** If  $X_r = A_1$ , noting that  $W = \{\text{id}, \sigma_\alpha\}$ , we have (2.7).

## § 6. Special Zeta-Values

Theorem 5.4 immediately implies the following theorem:

**Theorem 6.1** ([17]). For  $\mathbf{k} = (k_\alpha)_{\alpha \in \Delta_+} \in (2\mathbb{Z}_{\geq 1})^{|\Delta_+|}$  satisfying  $w^{-1}\mathbf{k} = \mathbf{k}$  for all  $w \in W$  (i.e.  $k_\alpha = k_\beta$  if  $\alpha$  and  $\beta$  are of the same length),

$$(6.1) \quad \zeta_r(\mathbf{k}, \mathbf{0}; \Delta) = \frac{(-1)^{|\Delta_+|}}{|W|} \mathcal{P}(\mathbf{k}, \mathbf{0}; \Delta) \left( \prod_{\alpha \in \Delta_+} \frac{(2\pi i)^{k_\alpha}}{k_\alpha!} \right) \in \mathbb{Q}\pi^{|\mathbf{k}|}$$

where  $|\mathbf{k}| = \sum_{\alpha \in \Delta_+} k_\alpha$ .

**Example 6.2.** If  $X_r = A_1$ , we have

$$(6.2) \quad \zeta(k) = \frac{-1}{2} B_k \frac{(2\pi i)^k}{k!} \in \mathbb{Q}\pi^k \quad (k \in 2\mathbb{Z}_{\geq 1}).$$

In particular,  $\mathbf{k} = (k)_{\alpha \in \Delta_+}$  with  $k \in 2\mathbb{Z}_{\geq 1}$  (that is, all  $k_\alpha = k$ ) satisfies the condition in Theorem 2. In this case,  $\zeta_r(\mathbf{k}, \mathbf{0}; \Delta) \in \mathbb{Q}\pi^{|\Delta_+|k}$  was shown by Witten and Zagier. In our method, the rational factor is explicitly evaluated via the generating function. Our statement is indeed a non-trivial generalization of their results since we also have for example,

$$(6.3) \quad \begin{aligned} \zeta_2((2, 4, 4, 2), \mathbf{0}; C_2) &= \sum_{m, n=1}^{\infty} \frac{1}{m^2 n^4 (m+n)^4 (m+2n)^2} \\ &= \frac{(-1)^4}{2^2 2!} \frac{53}{1513512000} \left( \frac{(2\pi i)^2}{2!} \right)^2 \left( \frac{(2\pi i)^4}{4!} \right)^2 \\ &= \frac{53\pi^{12}}{6810804000}. \end{aligned}$$

## § 7. Multiple Periodic Bernoulli Functions

In this section, we give the definition of generating functions of multiple periodic Bernoulli functions. Let  $\mathcal{V}$  be the set of all  $\mathbb{R}$ -bases  $\mathbf{V} \subset \Delta_+$  and let  $\mathbf{V}^\vee = \{\beta^\vee\}_{\beta \in \mathbf{V}}$ . Let



$\mathbf{V}^* = \{\mu_\beta^{\mathbf{V}}\}_{\beta \in \mathbf{V}}$  be the dual basis of  $\mathbf{V}^{\vee}$ . Let  $Q^{\vee} = \bigoplus_{i=1}^r \mathbb{Z}\alpha_i^{\vee}$  be the coroot lattice and  $L(\mathbf{V}^{\vee}) = \bigoplus_{\beta \in \mathbf{V}} \mathbb{Z}\beta^{\vee}$ , which is a sublattice of  $Q^{\vee}$  with finite index ( $|Q^{\vee}/L(\mathbf{V}^{\vee})| < \infty$ ).

Fix a certain  $\phi \in V$  and define a multiple generalization of the notion of “fractional part” of  $\mathbf{y} \in V$  as

$$(7.1) \quad \{\mathbf{y}\}_{\mathbf{v},\beta} = \begin{cases} \{\langle \mathbf{y}, \mu_\beta^{\mathbf{V}} \rangle\} & (\langle \phi, \mu_\beta^{\mathbf{V}} \rangle > 0), \\ 1 - \{-\langle \mathbf{y}, \mu_\beta^{\mathbf{V}} \rangle\} & (\langle \phi, \mu_\beta^{\mathbf{V}} \rangle < 0). \end{cases}$$

Using these definitions, we have

**Definition 7.1** (The generating function [12, 16, 17]). For  $\mathbf{t} = (t_\alpha)_{\alpha \in \Delta_+} \in \mathbb{C}^{|\Delta_+|}$ ,

$$(7.2) \quad F(\mathbf{t}, \mathbf{y}; \Delta) = \sum_{\mathbf{v} \in \mathcal{V}} \left( \prod_{\gamma \in \Delta_+ \setminus \mathbf{v}} \frac{t_\gamma}{t_\gamma - \sum_{\beta \in \mathbf{v}} t_\beta \langle \gamma^{\vee}, \mu_\beta^{\mathbf{V}} \rangle} \right) \times \frac{1}{|Q^{\vee}/L(\mathbf{V}^{\vee})|} \sum_{q \in Q^{\vee}/L(\mathbf{V}^{\vee})} \left( \prod_{\beta \in \mathbf{V}} \frac{t_\beta \exp(t_\beta \{\mathbf{y} + q\}_{\mathbf{v},\beta})}{e^{t_\beta} - 1} \right).$$

It can be shown that the generating function  $F(\mathbf{t}, \mathbf{y}; \Delta)$  is holomorphic in the neighborhood of the origin in  $\mathbf{t}$ .

**Definition 7.2** (Multiple periodic Bernoulli functions [12, 16, 17]). We define multiple periodic Bernoulli functions  $\mathcal{P}(\mathbf{k}, \mathbf{y}; \Delta)$  by the coefficients of the Taylor expansion

$$(7.3) \quad F(\mathbf{t}, \mathbf{y}; \Delta) = \sum_{\mathbf{k} \in \mathbb{Z}_{\geq 0}^{|\Delta_+|}} \mathcal{P}(\mathbf{k}, \mathbf{y}; \Delta) \prod_{\alpha \in \Delta_+} \frac{t_\alpha^{k_\alpha}}{k_\alpha!}.$$

**Example 7.3.** If  $X_r = A_1$ , we have

$$(7.4) \quad F(t, y) = \frac{te^{t\{y\}}}{e^t - 1} = \sum_{k=0}^{\infty} B_k(\{y\}) \frac{t^k}{k!}.$$

From this example, we see that  $\mathcal{P}(\mathbf{k}, \mathbf{y}; \Delta)$  can be regarded as natural generalizations of  $B_k(\{y\})$ .

## § 8. An Example: $A_2$ Case

We calculate a multiple periodic Bernoulli function and its generating function in the case of the root system of type  $A_2$ .


We have the basic data as follows:

$$\Delta_+^{\vee} = \{\alpha_1^{\vee}, \alpha_2^{\vee}, \alpha_1^{\vee} + \alpha_2^{\vee}\}, \quad \mathcal{V} = \{\mathbf{V}_1, \mathbf{V}_2, \mathbf{V}_3\},$$

$$\mathbf{t} = (t_{\alpha_1}, t_{\alpha_2}, t_{\alpha_1 + \alpha_2}) = (t_1, t_2, t_3),$$

$$\mathbf{y} = y_1 \alpha_1^{\vee} + y_2 \alpha_2^{\vee},$$

where  $\mathbf{V}_1 = \{\alpha_1, \alpha_2\}$ ,  $\mathbf{V}_2 = \{\alpha_1, \alpha_1 + \alpha_2\}$  and  $\mathbf{V}_3 = \{\alpha_2, \alpha_1 + \alpha_2\}$ . By definition, we also



have  $\mathbf{V}_1^\vee = \{\alpha_1^\vee, \alpha_2^\vee\}$ ,  $\mathbf{V}_2^\vee = \{\alpha_1^\vee, \alpha_1^\vee + \alpha_2^\vee\}$ ,  $\mathbf{V}_3^\vee = \{\alpha_2^\vee, \alpha_1^\vee + \alpha_2^\vee\}$  and  $\mathbf{V}_1^* = \{\lambda_1, \lambda_2\}$ ,  $\mathbf{V}_2^* = \{\lambda_1 - \lambda_2, \lambda_2\}$ ,  $\mathbf{V}_3^* = \{\lambda_2 - \lambda_1, \lambda_1\}$ . Fix a sufficiently small  $\varepsilon > 0$  and  $\phi = \alpha_1^\vee + \varepsilon\alpha_2^\vee$ . Then by Definition 7.1 and using these data, we have the generating function as

$$(8.1a) \quad F(\mathbf{t}, \mathbf{y}; A_2) = \frac{t_3}{t_3 - t_1 - t_2} \frac{t_1 e^{t_1 \{y_1\}}}{e^{t_1} - 1} \frac{t_2 e^{t_2 \{y_2\}}}{e^{t_2} - 1}$$

$$(8.1b) \quad + \frac{t_2}{t_2 + t_1 - t_3} \frac{t_1 e^{t_1 \{y_1 - y_2\}}}{e^{t_1} - 1} \frac{t_3 e^{t_3 \{y_2\}}}{e^{t_3} - 1}$$

$$(8.1c) \quad + \frac{t_1}{t_1 + t_2 - t_3} \frac{t_2 e^{t_2(1 - \{y_1 - y_2\})}}{e^{t_2} - 1} \frac{t_3 e^{t_3 \{y_1\}}}{e^{t_3} - 1},$$

where (8.1a), (8.1b) and (8.1c) correspond to  $\mathbf{V}_1$ ,  $\mathbf{V}_2$  and  $\mathbf{V}_3$  respectively. Note that in this case,  $L(\mathbf{V}_1^\vee) = L(\mathbf{V}_2^\vee) = L(\mathbf{V}_3^\vee) = Q^\vee$  and the second sum of (7.2) is trivial.

For  $\mathbf{k} = \mathbf{2} = (2, 2, 2)$ , expanding the right-hand side of (8.1a)–(8.1c), we find that the multiple periodic Bernoulli function is

$$(8.2) \quad \begin{aligned} \mathcal{P}(\mathbf{2}, (y_1, y_2); A_2) &= \frac{1}{3780} + \frac{1}{90}(\{y_1\} - \{y_1 - y_2\} - \{y_2\}) \\ &+ \frac{1}{90}(-\{y_1\}^2 - 2\{y_1 - y_2\}\{y_1\} + \{y_1 - y_2\}^2 - \{y_2\}^2 + 2\{y_1 - y_2\}\{y_2\}) \\ &+ \frac{1}{18}(-\{y_1\}^3 + 3\{y_1 - y_2\}\{y_1\}^2 + 3\{y_2\}^3 + 3\{y_1 - y_2\}\{y_2\}^2) \\ &+ \frac{1}{18}(\{y_1\}^4 - 2\{y_1 - y_2\}\{y_1\}^3 - 3\{y_1 - y_2\}^2\{y_1\}^2 \\ &\quad - 5\{y_2\}^4 - 10\{y_1 - y_2\}\{y_2\}^3 - 3\{y_1 - y_2\}^2\{y_2\}^2) \\ &+ \frac{1}{30}(\{y_1\}^5 - 5\{y_1 - y_2\}\{y_1\}^4 + 10\{y_1 - y_2\}^2\{y_1\}^3 \\ &\quad + 5\{y_2\}^5 + 15\{y_1 - y_2\}\{y_2\}^4 + 10\{y_1 - y_2\}^2\{y_2\}^3) \\ &+ \frac{1}{30}(-\{y_1\}^6 + 4\{y_1 - y_2\}\{y_1\}^5 - 5\{y_1 - y_2\}^2\{y_1\}^4 \\ &\quad - \{y_2\}^6 - 4\{y_1 - y_2\}\{y_2\}^5 - 5\{y_1 - y_2\}^2\{y_2\}^4). \end{aligned}$$

By Theorem 5.4, we have a functional relation in  $y_1, y_2$  corresponding to this multiple periodic Bernoulli function:

$$(8.3) \quad \begin{aligned} \zeta_2(\mathbf{2}, (y_1, y_2); A_2) &+ \zeta_2(\mathbf{2}, (-y_1 + y_2, y_2); A_2) + \zeta_2(\mathbf{2}, (y_1, y_1 - y_2); A_2) \\ &+ \zeta_2(\mathbf{2}, (-y_2, y_1 - y_2); A_2) + \zeta_2(\mathbf{2}, (-y_1 + y_2, -y_1); A_2) + \zeta_2(\mathbf{2}, (-y_2, -y_1); A_2) \\ &= (-1)^3 \mathcal{P}(\mathbf{2}, (y_1, y_2); A_2) \frac{(2\pi i)^6}{(2!)^3}. \end{aligned}$$

In particular if  $(y_1, y_2) = (0, 0)$ , then


$$(8.4) \quad \zeta_2(\mathbf{2}, (0, 0); A_2) = \frac{1}{6}(-1)^3 \frac{1}{3780} \frac{(2\pi i)^6}{(2!)^3} = \frac{\pi^6}{2835}.$$

**Example 8.1.** If  $X_r = A_1$ , we have

$$(8.5) \quad \zeta(2) = \frac{1}{2}(-1)^1 \frac{1}{6} \frac{(2\pi i)^2}{2!} = \frac{\pi^2}{6}, \quad B_2(\{y\}) = \frac{1}{6} - \{y\} + \{y\}^2.$$

### § 9. Multiple Bernoulli Polynomials

In the classical theory, Bernoulli polynomials can be derived by the analytic continuation of periodic Bernoulli functions. We explain this fact. Let  $\mathfrak{H} = \{y \in \mathbb{R} \mid \{y\} \in \mathbb{Z}\} = \mathbb{Z}$  (discontinuous points of  $\{y\}$ ). Let  $\mathbb{R} \setminus \mathfrak{H} = \coprod_{\nu \in \mathbb{Z}} \mathfrak{D}^{(\nu)}$ , where  $\mathfrak{D}^{(\nu)} = (\nu, \nu + 1)$ . From each  $\mathfrak{D}^{(\nu)}$  to  $\mathbb{C}$ , the function  $B(\{y\})$  is analytically continued to a polynomial function  $B_k^{(\nu)}(y) = B_k(y - \nu) \in \mathbb{Q}[y]$ .



A similar procedure works well in general cases and we can define multiple generalizations of Bernoulli polynomials. Let


$$(9.1) \quad \mathfrak{H} = \bigcup_{\mathbf{v} \in \mathcal{V}} \bigcup_{q \in Q^{\mathbf{v}}} \bigcup_{\beta \in \mathbf{V}} \{\mathbf{y} \in V \mid \{\mathbf{y} + q\}_{\mathbf{v}, \beta} \in \mathbb{Z}\}$$

(discontinuous points of  $\{\mathbf{y} + q\}_{\mathbf{v}, \beta}$  appearing in the generating function). Let

$$(9.2) \quad V \setminus \mathfrak{H} = \coprod_{\nu \in \mathfrak{J}} \mathfrak{D}^{(\nu)},$$

where  $\mathfrak{D}^{(\nu)}$  is an open connected component and  $\mathfrak{J}$  is a set of indices.


The above figure is the situation in the  $A_2$  case, where lines are  $\mathfrak{H}$  and open triangles are  $\mathfrak{D}^{(\nu)}$ . For example,  $\{\mathbf{y} + q\}_{\mathbf{v}_1, \alpha_1} = \{y_1 \alpha_1^{\vee} + y_2 \alpha_2^{\vee} + q, \lambda_1\} = \{y_1\} \in \mathbb{Z}$  gives the lines parallel to  $\alpha_2^{\vee}$ .



**Theorem 9.1** ([12, 16, 17]). From each region  $\mathfrak{D}^{(\nu)}$  to the whole space  $\mathbb{C} \otimes V$ ,  $\mathcal{P}(\mathbf{k}, \mathbf{y}; \Delta)$  is analytically continued in  $\mathbf{y}$  to a polynomial function  $\mathcal{B}_{\mathbf{k}}^{(\nu)}(\mathbf{y}; \Delta) \in \mathbb{Q}[\mathbf{y}]$  of total degree at most  $|\mathbf{k}|$ , where  $\mathbf{y} = \sum_{n=1}^r y_n \alpha_n^{\vee}$ .

#### § 9.1. An Example: $A_2$ Case

The Bernoulli polynomial  $\mathcal{B}_2^{(0)}(\mathbf{y}; A_2)$  is obtained by the analytic continuation of the periodic Bernoulli function  $\mathcal{P}(\mathbf{2}, \mathbf{y}; A_2)$  from the region  $\mathfrak{D}^{(0)}$ , which is the shaded triangle region in the figure below.



$$V \setminus \mathfrak{H} = \coprod_{\nu \in \mathfrak{J}} \mathfrak{D}^{(\nu)}$$

$$\mathcal{P}(\mathbf{2}, \mathbf{y}; A_2)$$

$$\mathcal{B}_2^{(0)}(\mathbf{y}; A_2)$$

(Periodic Bernoulli function)


(Bernoulli polynomial)


The explicit form of the Bernoulli polynomial  $\mathcal{B}_2^{(0)}(\mathbf{y}; A_2)$  is given, simply by removing all curly brackets from (8.2), as follows:

$$\begin{aligned}
 (9.3) \quad \mathcal{B}_2^{(0)}(\mathbf{y}; A_2) &= \frac{1}{3780} + \frac{1}{45}(y_1 y_2 - y_1^2 - y_2^2) + \frac{1}{18}(3y_1 y_2^2 - 3y_1^2 y_2 + 2y_1^3) \\
 &\quad + \frac{1}{9}(-2y_1 y_2^3 - 3y_1^2 y_2^2 + 4y_1^3 y_2 - 2y_1^4 + y_2^4) \\
 &\quad + \frac{1}{30}(-5y_1 y_2^4 + 10y_1^2 y_2^3 + 10y_1^3 y_2^2 - 15y_1^4 y_2 + 6y_1^5) \\
 &\quad + \frac{1}{30}(6y_1 y_2^5 - 5y_1^2 y_2^4 - 5y_1^4 y_2^2 + 6y_1^5 y_2 - 2y_1^6 - 2y_2^6) \in \mathbb{Q}[\mathbf{y}].
 \end{aligned}$$

### § 9.2. Further Examples: $C_2, G_2$ Cases

The following graphs in the upper (resp. lower) row are of type  $C_2$  (resp.  $G_2$ ).





We summarize what we have obtained: We have constructed periodic Bernoulli functions so that they describe functional-relations in  $\mathbf{y}$  of multiple zeta-functions of root systems, which can be calculated by use of the generating function; Bernoulli polynomials are obtained by the analytic continuation of periodic Bernoulli functions.

### § 10. $L$ -Functions of Root Systems

We give another application of periodic Bernoulli functions or equivalently Bernoulli polynomials. For this purpose, we define an  $L$ -analogue of zeta-functions of root systems.

**Definition 10.1** ( $L$ -functions of root systems [12, 16]). For a root system  $\Delta$ , define

$$(10.1) \quad L_r(\mathbf{s}, \boldsymbol{\chi}; \Delta) = \sum_{\lambda \in P_+} \prod_{\alpha \in \Delta_+} \frac{\chi_\alpha(\langle \alpha^\vee, \lambda + \rho \rangle)}{\langle \alpha^\vee, \lambda + \rho \rangle^{s_\alpha}},$$

where  $\boldsymbol{\chi} = (\chi_\alpha)_{\alpha \in \Delta_+}$  is a set of primitive Dirichlet characters of conductors  $f_\alpha \in \mathbb{Z}_{\geq 1}$ .

We extend  $\boldsymbol{\chi} = (\chi_\alpha)_{\alpha \in \Delta_+}$  to  $(\chi_\alpha)_{\alpha \in \Delta}$  by  $\chi_\alpha = \chi_{-\alpha}$  and define  $(w\boldsymbol{\chi})_\alpha = \chi_{w^{-1}\alpha}$ . Then we have value-relations of  $L$ -functions.

**Theorem 10.2** ([12, 16]). For  $\mathbf{s} = \mathbf{k} = (k_\alpha)_{\alpha \in \Delta_+} \in \mathbb{Z}_{\geq 2}^{|\Delta_+|}$ , we have

$$(10.2) \quad \sum_{w \in W} \left( \prod_{\alpha \in \Delta_+ \cap w\Delta_-} (-1)^{k_\alpha} \chi_\alpha(-1) \right) L_r(w^{-1}\mathbf{k}, w^{-1}\boldsymbol{\chi}; \Delta) \\ = (-1)^{|\Delta_+|} \left( \prod_{\alpha \in \Delta_+} \chi_\alpha(-1) g(\chi_\alpha) \frac{(2\pi i)^{k_\alpha}}{k_\alpha! f^{k_\alpha}} \right) \mathcal{B}_{\mathbf{k}, \bar{\boldsymbol{\chi}}}(\Delta),$$

where  $\mathcal{B}_{\mathbf{k}, \bar{\boldsymbol{\chi}}}(\Delta)$  is a multiple generalized Bernoulli number (defined later).

**Example 10.3.** If  $X_r = A_1$ , we have the classical result

$$(10.3) \quad L(k, \chi) + (-1)^k \chi(-1) L(k, \chi) = -\chi(-1) g(\chi) \frac{(2\pi i)^k}{k! f^k} B_{k, \bar{\chi}},$$

where  $B_{k, \bar{\chi}}$  is the generalized Bernoulli number given in (2.10). As for the traditional account of this formula, see [3, chapter 1] for example.

### § 11. Special $L$ -Values

Theorem 10.2 immediately implies a formula for special values of  $L$ -functions:

**Theorem 11.1** ([12, 16]). For  $\mathbf{k} \in (\mathbb{Z}_{\geq 2})^{|\Delta_+|}$  and  $\chi$  such that  $w^{-1}\mathbf{k} = \mathbf{k}$ ,  $w^{-1}\chi = \chi$  for all  $w \in W$  and  $(-1)^{k_\alpha} \chi_\alpha(-1) = 1$  for all  $\alpha \in \Delta_+$ , we have

$$(11.1) \quad L_r(\mathbf{k}, \chi; \Delta) = \frac{(-1)^{|\mathbf{k}|+|\Delta_+|}}{|W|} \left( \prod_{\alpha \in \Delta_+} \frac{(2\pi i)^{k_\alpha}}{k_\alpha! f_\alpha^{k_\alpha}} g(\chi_\alpha) \right) \mathcal{B}_{\mathbf{k}, \bar{\chi}}(\Delta).$$

**Example 11.2.** If  $X_r = A_1$ , we have

$$(11.2) \quad L(k, \chi) = \frac{(-1)^{k+1}}{2} \frac{(2\pi i)^k}{k! f^k} g(\chi) B_{k, \bar{\chi}}.$$

**Example 11.3.** Let  $\rho_7$  be the Dirichlet character of conductor 7 defined by  $\rho_7(1) = \rho_7(6) = 1$ ,  $\rho_7(2) = \rho_7(5) = e^{2\pi i/3}$ ,  $\rho_7(3) = \rho_7(4) = e^{4\pi i/3}$ . Then the associated Gauss sum is  $g(\rho_7) = 2(\cos(2\pi/7) + e^{2\pi i/3} \cos(4\pi/7) + e^{4\pi i/3} \cos(6\pi/7))$  and we have

$$(11.3) \quad L_2((2, 4, 4, 2), (1, \rho_7, \rho_7, 1); C_2) = \sum_{m, n=1}^{\infty} \frac{\rho_7(n) \rho_7(m+n)}{m^2 n^4 (m+n)^4 (m+2n)^2} \\ = \frac{(-1)^{12+4}}{2^2 2!} \left( \frac{(2\pi i)^2}{2!} \right)^2 \left( \frac{(2\pi i)^4}{4! 7^4} g(\rho_7) \right)^2 \left( \frac{69967019}{6988350600} + \frac{102810289\sqrt{-3}}{6988350600} \right) \\ = g(\rho_7)^2 \pi^{12} \left( \frac{69967019}{181289027372537700} + \frac{102810289\sqrt{-3}}{181289027372537700} \right).$$

**Example 11.4.** Let  $\rho_5$  be the quadratic character of conductor 5 given in Theorem B. Then we have

$$(11.4) \quad L_2((2, 2, 2, 2), (\rho_5, \rho_5, \rho_5, \rho_5); C_2) = \frac{92}{29296875} \pi^8;$$

$$(11.5) \quad L_3((2, 2, 2, 2, 2, 2), (\rho_5, \rho_5, \rho_5, \rho_5, \rho_5, \rho_5); A_3) = -\frac{1856}{213623046875} \pi^{12}.$$

The latter can be regarded as a character analogue of the formula in [6, Prop. 8.5].

## § 12. Multiple Generalized Bernoulli Numbers

The generating function of multiple generalized Bernoulli numbers is given in terms of that of multiple Bernoulli polynomials as in the classical theory.

**Definition 12.1** (The generating function [12, 16]). For  $\mathbf{t} = (t_\alpha)_{\alpha \in \Delta_+}$ ,

$$(12.1) \quad G(\mathbf{t}, \chi; \Delta) = \sum_{\substack{a_\alpha=1 \\ \alpha \in \Delta_+}}^{f_\alpha} \left( \prod_{\alpha \in \Delta_+} \frac{\chi_\alpha(a_\alpha)}{f_\alpha} \right) F(\mathbf{f} \mathbf{t}, \mathbf{y}(\mathbf{a}; \mathbf{f}); \Delta),$$

where  $F(\mathbf{t}, \mathbf{y}; \Delta)$  is the generating function of multiple periodic Bernoulli functions in Definition 7.1 and  $\mathbf{f} \mathbf{t} = (f_\alpha t_\alpha)_{\alpha \in \Delta_+}$ ,  $\mathbf{y}(\mathbf{a}; \mathbf{f}) = \sum_{\alpha \in \Delta_+} a_\alpha \alpha^\vee / f_\alpha$ .

**Definition 12.2** (Multiple generalized Bernoulli numbers [12, 16]). We define multiple generalized Bernoulli numbers  $\mathcal{B}_{\mathbf{k}, \chi}(\Delta)$  by the coefficients of the Taylor expansion

$$(12.2) \quad G(\mathbf{t}, \chi; \Delta) = \sum_{\mathbf{k} \in \mathbb{Z}_{\geq 0}^{|\Delta_+|}} \mathcal{B}_{\mathbf{k}, \chi}(\Delta) \prod_{\alpha \in \Delta_+} \frac{t_\alpha^{k_\alpha}}{k_\alpha!}.$$

We note that  $\mathcal{B}_{\mathbf{k}, \chi}(\Delta)$  can be written in terms of multiple periodic Bernoulli functions as

$$(12.3) \quad \mathcal{B}_{\mathbf{k}, \chi}(\Delta) = \left( \prod_{\alpha \in \Delta_+} f_\alpha^{k_\alpha - 1} \right) \sum_{\substack{a_\alpha=1 \\ \alpha \in \Delta_+}}^{f_\alpha} \left( \prod_{\alpha \in \Delta_+} \chi_\alpha(a_\alpha) \right) \mathcal{P}(\mathbf{k}, \mathbf{y}(\mathbf{a}; \mathbf{f}); \Delta).$$

**Example 12.3.** If  $X_r = A_1$ , we have the generating function

$$(12.4) \quad G(t, \chi) = \sum_{a=1}^f \frac{\chi(a)}{f} F(ft, a/f) = \sum_{a=1}^f \frac{\chi(a)}{f} \frac{f t e^{ft\{a/f\}}}{e^{ft} - 1} = \sum_{k=0}^{\infty} B_{k, \chi} \frac{t^k}{k!}.$$

**Theorem 12.4** ([12, 16]). Assume that  $\Delta$  is irreducible. Moreover assume that  $f_\alpha > 1$  if  $\Delta$  is of type  $A_1$ . Then for  $w \in W$ , we have

$$(12.5) \quad B_{w^{-1}\mathbf{k}, w^{-1}\chi}(\Delta) = \left( \prod_{\alpha \in \Delta_+ \cap w\Delta_-} (-1)^{k_\alpha} \chi_\alpha(-1) \right) \mathcal{B}_{\mathbf{k}, \chi}(\Delta).$$

Hence  $\mathcal{B}_{\mathbf{k}, \chi}(\Delta) = 0$  if there exists an element  $w \in W_{\mathbf{k}} \cap W_\chi$  such that

$$(12.6) \quad \prod_{\alpha \in \Delta_+ \cap w\Delta_-} (-1)^{k_\alpha} \chi_\alpha(-1) \neq 1,$$

where  $W_{\mathbf{k}}$  and  $W_\chi$  are the stabilizers of  $\mathbf{k}$  and  $\chi$  respectively.

**Example 12.5.** If  $X_r = A_1$ , we have

$$(12.7) \quad B_{k,\chi} = 0 \quad \text{if } (-1)^k \chi(-1) \neq 1.$$

Several other properties in the classical theory such as

$$F(t, y) = F(-t, -y) \text{ for } y \in \mathbb{R} \setminus \mathbb{Z}, \quad B_k(1 - y) = (-1)^k B_k(y), \quad \frac{1}{t} \frac{\partial}{\partial y} F(t, y) = F(t, y)$$

can be reinterpreted in terms of root systems and Weyl groups.

### § 13. Zeta-Functions for Lie Groups

Recall that volume formulas are associated with all connected compact semisimple Lie groups. It is known that there is one-to-one correspondence between finite dimensional representations of complex semisimple Lie algebra  $\mathfrak{g}$  and those of connected simply-connected compact semisimple Lie group  $G$ . In the cases of general compact semisimple Lie groups, we need analytically integral forms  $L$  for a maximal torus of  $G$ , which satisfies  $Q \subset L \subset P$ .

**Definition 13.1** (Zeta-functions of Lie groups). For a connected compact semisimple Lie group  $G$ ,

$$(13.1) \quad \zeta_r(\mathbf{s}, \mathbf{y}; G) = \sum_{\lambda \in L \cap P_+} e^{2\pi i \langle \mathbf{y}, \lambda + \rho \rangle} \prod_{\alpha \in \Delta_+} \frac{1}{\langle \alpha^\vee, \lambda + \rho \rangle^{s_\alpha}}.$$

**Lemma 13.2.**

$$(13.2) \quad \zeta_r(\mathbf{s}, \mathbf{y}; G) = \sum_{\mu \in P^\vee / Q^\vee} \widehat{\iota_{L+\rho}}(\mu) \zeta_r(\mathbf{s}, \mathbf{y} + \mu; \Delta),$$

where  $\widehat{\iota_{L+\rho}} : P^\vee / Q^\vee \rightarrow \mathbb{C}$  is the Fourier transformation of the characteristic function of  $L + \rho$  given by

$$(13.3) \quad \widehat{\iota_{L+\rho}}(\mu) = \frac{1}{|P/Q|} \sum_{\lambda \in (L+\rho)/Q} e^{-2\pi i \langle \mu, \lambda \rangle}.$$

Note that this expression plays the same role as the finite Fourier transformation of the Dirichlet character (see [35, Lemma 4.7]) in the theory of Dirichlet  $L$ -functions, whose origin is the study of prime numbers satisfying congruence conditions. In fact, our  $\zeta_r(\mathbf{s}, \mathbf{y}; G)$  is a kind of Dirichlet series with congruence conditions (see (13.8) as an example).

In the  $A_1$  case with  $L = Q$ , Lemma 13.2 implies

$$(13.4) \quad \sum_{m=0}^{\infty} \frac{e^{2\pi i(2m+1)y}}{(2m+1)^s} = \sum_{m=0}^{\infty} \frac{1}{2} \frac{e^{2\pi i(m+1)y}}{(m+1)^s} + \sum_{m=0}^{\infty} \frac{-1}{2} \frac{e^{2\pi i(m+1)(y+\frac{1}{2})}}{(m+1)^s}.$$



**Lemma 13.3.** For  $\mu \in P^\vee/Q^\vee$ , we have

$$(13.5) \quad \widehat{\iota_{L+\rho}}(\mu) = \frac{(-1)^{\langle \mu, 2\rho \rangle}}{|\pi_1(G)|} \delta_{L^*/Q^\vee}(\mu) \in \frac{\{-1, 0, 1\}}{|\pi_1(G)|} \subset \mathbb{Q},$$

where  $\pi_1(G)$  denotes the fundamental group of  $G$  and

$$(13.6) \quad \delta_{L^*/Q^\vee}(\mu) = \begin{cases} 1 & (\mu \in L^*/Q^\vee), \\ 0 & (\mu \notin L^*/Q^\vee). \end{cases}$$

Noting  $P/L \simeq L^*/Q^\vee \simeq \pi_1(G)$ , we have the following, where  $G$  may not be simply-connected.

**Theorem 13.4** ([22]). For  $\mathbf{k} = (k_\alpha)_{\alpha \in \Delta_+} \in (2\mathbb{Z}_{\geq 1})^{|\Delta_+|}$  satisfying  $w^{-1}\mathbf{k} = \mathbf{k}$  for all  $w \in W$ , and  $\nu \in P^\vee/Q^\vee$  (a central element of  $G$ ), we have

$$(13.7) \quad \zeta_r(\mathbf{k}, \nu; G) = \frac{(-1)^{|\Delta_+|}}{|W|} \mathcal{P}(\mathbf{k}, \nu; G) \left( \prod_{\alpha \in \Delta_+} \frac{(2\pi i)^{k_\alpha}}{k_\alpha!} \right) \in \mathbb{Q}\pi^{|\mathbf{k}|}.$$

As an example, we obtain for the projective unitary group  $PU(3)$ ,

$$(13.8) \quad \begin{aligned} \zeta_2(\mathbf{2}, \mathbf{0}; PU(3)) &= \sum_{\substack{m, n=1 \\ m \equiv n \pmod{3}}}^{\infty} \frac{1}{m^2 n^2 (m+n)^2} \\ &= \sum_{2m-n, 2n-m > 0} \frac{1}{(2m-n)^2 (2n-m)^2 (m+n)^2} \\ &= \frac{(-1)^3}{3!} \frac{187}{918540} \left( \frac{(2\pi i)^2}{2!} \right)^3 \\ &= \frac{187\pi^6}{688905}. \end{aligned}$$

*Remark.* Originally, Witten zeta-functions represent the volumes of certain moduli spaces. Introducing multi-variable generalizations, we find some new applications. For example, we give a new interpretation of the shuffle product in the theory of Euler-Zagier multiple zeta values [19] and evaluate a class of Euler-Zagier multiple zeta values [20, 23]. However the geometric meaning of special values of zeta-functions of root systems is yet to be clarified.

## § 14. An Integral Representation

This section is based on the results by the first author in [9, 10]. So far, we focused on special values on the region of convergence. On the other hand, analytic continuations enable us to discuss special values on the whole space in  $\mathbf{s}$ .

The analytic continuations of general multiple zeta-functions were already obtained by Lichtin [24], Essouabri [4, 5], Matsumoto [25, 26], de Crisenoy [2], etc. (See [27] for an elaborated survey on the analytic continuations of multiple zeta-functions.) However we give yet another method which is a generalization of the formula

$$(14.1) \quad \zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \frac{1}{\Gamma(s)(e^{2\pi i s} - 1)} \int_C \frac{z^{s-1}}{e^z - 1} dz \quad (C: \text{Hankel contour}).$$

Let  $N, R$  be positive integers. For  $\boldsymbol{\xi} = (\xi_1, \dots, \xi_R) \in \mathbb{C}^R$ ,  $\mathbf{a} = (a_1, \dots, a_N)$ ,  $\mathbf{s} = (s_1, \dots, s_N) \in \mathbb{C}^N$  and  $\mathbf{b} = (b_{ij})_{1 \leq i \leq N, 1 \leq j \leq R} \in \mathbb{C}^{N \times R}$ , consider the multiple series

$$(14.2) \quad \zeta(\boldsymbol{\xi}, \mathbf{a}, \mathbf{b}, \mathbf{s}) = \sum_{m_1=0}^{\infty} \cdots \sum_{m_R=0}^{\infty} \frac{e^{\xi_1 m_1} \cdots e^{\xi_R m_R}}{(a_1 + b_{11}m_1 + \cdots + b_{1R}m_R)^{s_1} \cdots (a_N + b_{N1}m_1 + \cdots + b_{NR}m_R)^{s_N}}.$$

**Theorem 14.1** ([9, 10]).

$$(14.3) \quad \zeta(\boldsymbol{\xi}, \mathbf{a}, \mathbf{b}, \mathbf{s}) = \frac{1}{\Gamma(s_1) \cdots \Gamma(s_N)} \prod_{t \in S} \frac{1}{e^{2\pi i t(\mathbf{s})} - 1} \times \int_{\Sigma} \frac{e^{(b_{11} + \cdots + b_{1R} - a_1)z_1} \cdots e^{(b_{N1} + \cdots + b_{NR} - a_N)z_N} z_1^{s_1-1} \cdots z_N^{s_N-1}}{(e^{z_1 b_{11} + \cdots + z_N b_{N1}} - e^{\xi_1}) \cdots (e^{z_1 b_{1R} + \cdots + z_N b_{NR}} - e^{\xi_R})} dz_1 \wedge \cdots \wedge dz_N,$$

where  $\Sigma$  is a union of certain surfaces and  $S$  is a set of certain linear functionals on  $\mathbb{C}^N$ .

If  $b_{ij} > 0$  for all  $i, j$  satisfying  $1 \leq i \leq N$ ,  $1 \leq j \leq R$ , then this integral representation can be derived by use of Shintani's result [30]. In fact, Theorem 14.1 is a refinement of his integral representation.

Setting  $\xi_i = 0$ ,  $a_{\alpha} = \langle \alpha^{\vee}, \rho \rangle$  and  $b_{\alpha i} = \langle \alpha^{\vee}, \lambda_i \rangle$  for  $\alpha \in \Delta_+$  and  $1 \leq i \leq R = r$ , we obtain integral representations of zeta-functions of root systems. In this setting, from the integrand, we can construct generating functions of Bernoulli numbers for nonpositive domain.

## § 15. Possibilities of Elliptic Generalizations

Lastly we give two possibilities of “elliptic” generalizations by regarding  $\zeta_r(\mathbf{s}, \mathbf{y}; \Delta)$  as “rational” or “trigonometric” versions.

The first is an Eisenstein analogue. Let  $k > 2$  be an integer,  $(x, y) \in \mathbb{R}^2 \setminus \mathbb{Z}^2$  and  $\tau \in \mathbb{C}$  with  $\Im \tau > 0$ . The Eisenstein series is defined by

$$(15.1) \quad G_k(\tau; x, y) = \sum_{(m,n) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \frac{e^{2\pi i(mx+ny)}}{(m+n\tau)^k}.$$

We define  $\mathcal{H}_k(x, y; \tau)$  by

$$(15.2) \quad e^{2\pi i x t} \frac{\theta'(0; \tau) \theta(t + x\tau - y; \tau)}{\theta(t; \tau) \theta(x\tau - y; \tau)} = \sum_{k=0}^{\infty} \mathcal{H}_k(x, y; \tau) \frac{(2\pi i)^k t^{k-1}}{k!},$$

where  $t \in \mathbb{C}$  with  $|t| < \epsilon$  for sufficiently small  $\epsilon > 0$  and  $\theta(z; \tau)$  is the Jacobi theta function defined by

$$(15.3) \quad \theta(z; \tau) = -i \sum_{n \in \mathbb{Z}} \exp\left(\pi i \left(n + \frac{1}{2}\right)^2 \tau + 2\pi i \left(n + \frac{1}{2}\right) z + \pi i n\right)$$

for  $z \in \mathbb{C}$ . Then we have the following, which can be regarded as an elliptic analogue of the result on the zeta-function of root system of type  $A_1$  given in (2.7).

**Proposition 15.1** (Katayama [8]). *For  $k \in \mathbb{Z}_{\geq 2}$ , we have*

$$(15.4) \quad G_k(\tau; x, y) = -\mathcal{H}_k(x, y; \tau) \frac{(2\pi i)^k}{k!}.$$

From this viewpoint, it is desirable to develop a theory on elliptic analogues of the results on zeta-functions of root systems mentioned in the previous sections, by constructing corresponding Eisenstein series. For example, we hope to extend (15.4) to that associated with root systems.

The second is a  $q$ -analogue. Instead of Weyl's dimension formula, we employ the character formula. For  $q = e^{-2\pi i/\tau}$ ,  $s, z \in \mathbb{C}$  with  $\Re z > 0$  and  $x \in \mathbb{R}$ , define

$$(15.5) \quad \zeta_q(s, z; x) = \sum_{m=1}^{\infty} \frac{e^{2\pi i m x} q^{mz}}{[m]_q^s}, \quad [m]_q = \frac{1 - q^m}{1 - q}, \quad [m]_q! = [m]_q [m-1]_q \cdots [1]_q.$$

Let

$$(15.6) \quad \psi(t) = \frac{\tau}{2\pi i} \frac{e^{2\pi i t/\tau} - 1}{e^{2\pi i t z/\tau}} = t + O(t^2)$$

be a local coordinate around the origin. Define  $\mathcal{Q}_k(x, y, z; \tau)$  by

$$(15.7) \quad e^{2\pi i x t} \frac{\theta'(0; \tau) \theta(t + x\tau - y; \tau)}{\theta(t; \tau) \theta(x\tau - y; \tau)} = \sum_{k=0}^{\infty} \mathcal{Q}_k(x, y, z; \tau) \left(\frac{2\pi i/\tau}{1 - q}\right)^k \frac{\psi'(t) \psi(t)^{k-1}}{[k]_q!}.$$

Then

**Theorem 15.2.** *For  $k \in \mathbb{N}$ ,  $0 < z < 1$  and  $x, y \in \mathbb{R}$  with  $y + kz \in \mathbb{Z}$ , we have*

$$(15.8) \quad \zeta_q(k, k(1 - z); x) + (-1)^k \zeta_q(k, kz; -x) = -\mathcal{Q}_k(x, y, z; \tau) \frac{1}{[k]_q!}.$$

This is a  $q$ -analogue of (2.7). Not only the result, but also the proof can be done analogously. In fact, formula (2.7) can be shown by a residue calculus on the space  $\mathbb{C}$ . Similarly, we can prove Theorem 15.2 employing the space  $\mathbb{C}/\tau\mathbb{Z}$ .

In particular, from this formula, we have for  $\tau = i$ ,

$$(15.9) \quad \zeta_q(2, 1; 0) = (1 - e^{-2\pi})^2 \frac{\pi - 3}{24\pi}, \quad \zeta_q(4, 2; 0) = (1 - e^{-2\pi})^4 \frac{30\pi^3 - 11\pi^4 + 3\varpi^4}{1440\pi^4},$$

where  $\varpi$  is the lemniscate constant defined by

$$(15.10) \quad \varpi = 2 \int_0^1 \frac{dx}{\sqrt{1-x^4}}.$$

By aid of generalizations of Eisenstein series, these special values are also calculated in [21].

We hope that generalizations of the above will be constructed in arbitrary root systems.

## References

- [1] D. J. Broadhurst, J. M. Borwein, and D. M. Bradley, *Evaluation of  $k$ -fold Euler/Zagier sums: a compendium of results for arbitrary  $k$* , Electron. J. Combin. **4(2)** (1997), Research Paper 5, approx. 21 pp.
- [2] M. de Crisenoy, *Values at  $T$ -tuples of negative integers of twisted multivariable zeta series associated to polynomials of several variables*, Compos. Math. **142** (2006), 1373–1402.
- [3] K. Dilcher, *Zeros of Bernoulli, generalized Bernoulli and Euler polynomials*, Memoirs of Amer. Math. Soc. **386** (1988).
- [4] D. Essouabri, *Singularités des séries de Dirichlet associées à des polynômes de plusieurs variables et applications à la théorie analytique des nombres*, Thèse, Univ. Nancy I, 1995.
- [5] D. Essouabri, *Singularité des séries de Dirichlet associées à des polynômes de plusieurs variables et applications en théorie analytique des nombres*, Ann. Inst. Fourier (Grenoble) **47** (1997), 429–483.
- [6] P. E. Gunnells and R. Sczech, *Evaluation of Dedekind sums, Eisenstein cocycles, and special values of  $L$ -functions*, Duke Math. J. **118** (2003), 229–260.
- [7] J. G. Huard, K. S. Williams and N.-Y. Zhang, *On Tornheim’s double series*, Acta Arith. **75** (1996), 105–117.
- [8] K. Katayama, *On the values of Eisenstein series*, Tokyo J. Math. **1** (1978), 157–188.
- [9] Y. Komori, *An integral representation of Mordell-Tornheim double zeta function and its values at non-positive integers*, Ramanujan J. **17** (2008), 163–183.
- [10] Y. Komori, *An integral representation of multiple Hurwitz-Lerch zeta functions and generalized multiple Bernoulli numbers*, Quart. J. Math. **61** (2010), 437–496.
- [11] Y. Komori, K. Matsumoto and H. Tsumura, *Zeta-functions of root systems*, in “Proceedings of the Conference on  $L$ -functions” (Fukuoka, 2006), L. Weng and M. Kaneko (eds.), World Scientific, 2007, pp. 115–140.
- [12] Y. Komori, K. Matsumoto and H. Tsumura, *Zeta and  $L$ -functions and Bernoulli polynomials of root systems*, Proc. Japan Acad. Ser. A Math. Sci. **84** (2008), 57–62.

- [13] Y. Komori, K. Matsumoto and H. Tsumura, *On multiple Bernoulli polynomials and multiple  $L$ -functions of root systems*, in ‘Analytic Number Theory and Related Areas’, RIMS Kôkyûroku **1665**, Kyoto Univ., 2009, pp. 114–126.
- [14] Y. Komori, K. Matsumoto and H. Tsumura, *On multiple Bernoulli polynomials and multiple zeta-functions of root systems*, in ‘Representation Theory and Combinatorics’, RIMS Kôkyûroku **1689**, Kyoto Univ., 2010, pp. 117–132.
- [15] Y. Komori, K. Matsumoto and H. Tsumura, *On Witten multiple zeta-functions associated with semisimple Lie algebras II*, J. Math. Soc. Japan **62** (2010), 355–394.
- [16] Y. Komori, K. Matsumoto and H. Tsumura, *On multiple Bernoulli polynomials and multiple  $L$ -functions of root systems*, Proc. London Math. Soc. **100** (2010), 303–347.
- [17] Y. Komori, K. Matsumoto and H. Tsumura, *On Witten multiple zeta-functions associated with semisimple Lie algebras III*, to appear in “The Proceedings of the Conference on Multiple Dirichlet Series and Applications to Automorphic Forms” (Edinburgh, 2008), D. Bump et al. (eds.), arXiv:0907.0955.
- [18] Y. Komori, K. Matsumoto and H. Tsumura, *On Witten multiple zeta-functions associated with semisimple Lie algebras IV*, Glasg. Math. J. **53** (2011), 185–206.
- [19] Y. Komori, K. Matsumoto and H. Tsumura, *Shuffle products for multiple zeta values and partial fraction decompositions of zeta-functions of root systems*, to appear in Math. Z.
- [20] Y. Komori, K. Matsumoto and H. Tsumura, *Zeta-functions of root systems and multiple zeta values*, to appear in RIMS Kôkyûroku (in Japanese).
- [21] Y. Komori, K. Matsumoto and H. Tsumura, *Hyperbolic-sine analogues of Eisenstein series, generalized Hurwitz numbers, and  $q$ -zeta functions*, preprint, submitted for publication, arXiv:1006.3339.
- [22] Y. Komori, K. Matsumoto and H. Tsumura, *Zeta-functions of weight lattices of compact semisimple connected Lie groups*, preprint, submitted for publication, arXiv:1011.0323.
- [23] Y. Komori, K. Matsumoto and H. Tsumura, *Multiple zeta values and zeta-functions of root systems*, preprint, submitted for publication.
- [24] B. Lichtin, *The asymptotics of a lattice point problem associated to a finite number of polynomials*, Duke Math. J. **63** (1991), 139–192.
- [25] K. Matsumoto, *On Mordell-Tornheim and other multiple zeta-functions*, in “Proc. Session in Analytic Number Theory and Diophantine Equations”, D. R. Heath-Brown and B. Z. Moroz (eds.), Bonner Math. Schriften **360**, Bonn, 2003, n. 25, 17pp.
- [26] K. Matsumoto, *Analytic properties of multiple zeta-functions in several variables*, in “Number Theory: Tradition and Modernization”, Proc. 3rd China-Japan Seminar, W. Zhang and Y. Tanigawa (eds.), Springer, 2006, pp. 153–173.
- [27] K. Matsumoto, *The analytic theory of multiple zeta-functions and its applications*, Sugaku **59** (2007), 24–45 (in Japanese); English Transl.: Sugaku Expositions, **23** (2010), 143–168.
- [28] K. Matsumoto and H. Tsumura, *On Witten multiple zeta-functions associated with semisimple Lie algebras I*, Ann. Inst. Fourier (Grenoble) **56** (2006), 1457–1504.
- [29] L. J. Mordell, *On the evaluation of some multiple series*, J. London Math. Soc. **33** (1958), 368–371.
- [30] T. Shintani, *On evaluation of zeta functions of totally real algebraic number fields at non-positive integers*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **23** (1976), 393–417.
- [31] M. V. Subbarao and R. Sitaramachandrarao, *On some infinite series of L. J. Mordell and their analogues*, Pacific J. Math. **119** (1985), 245–255.
- [32] A. Szenes, *Iterated residues and multiple Bernoulli polynomials*, Internat. Math. Res. Notices **18** (1998), 937–958.

- [33] L. Tornheim, *Harmonic double series*, Amer. J. Math. **72** (1950), 303–314.
- [34] H. Tsumura, *On some combinatorial relations for Tornheim's double series*, Acta Arith. **105** (2002), 239–252.
- [35] L. C. Washington, *Introduction to Cyclotomic Fields*, 2nd ed., Springer-Verlag, New York Berlin Heidelberg, 1997.
- [36] E. Witten, *On quantum gauge theories in two dimensions*, Comm. Math. Phys. **141** (1991), 153–209.
- [37] D. Zagier, *Values of zeta functions and their applications*, in “First European Congress of Mathematics” Vol. II, A. Joseph et al. (eds.), Progr. Math. **120**, Birkhäuser, 1994, pp. 497–512.